


10049957 021502  
JC07 Rec'd PCT/PTO 19 FEB 2002

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER <b>KATO=21</b>
<b>TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371</b>		U.S. APPLICATION NO. (If known, see 37 CFR 1.5) <b>10/049957</b>
INTERNATIONAL APPLICATION NO. <b>PCT/JP00/05590</b>	INTERNATIONAL FILING DATE <b>21 August 2000</b>	PRIORITY CLAIMED <b>19 August 1999</b>
TITLE OF INVENTION <b>CHONDROGENESIS PROMOTERS</b>		
APPLICANT(S) FOR DO/EO/US <b>Yukio KATO et al.</b>		
<p>Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information.</p> <ol style="list-style-type: none"> <li><input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371</li> <li><input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371.</li> <li><input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).</li> <li><input checked="" type="checkbox"/> The US has been elected in a Demand by the expiration of 19 months from the priority date (PCT Article 31).</li> <li><input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2))             <ol style="list-style-type: none"> <li><input type="checkbox"/> is attached hereto (required only if not transmitted by the International Bureau)</li> <li><input checked="" type="checkbox"/> has been communicated by the International Bureau.</li> <li><input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US)</li> </ol> </li> <li><input checked="" type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2))</li> <li><input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))             <ol style="list-style-type: none"> <li><input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau).</li> <li><input type="checkbox"/> have been communicated by the International Bureau.</li> <li><input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</li> <li><input checked="" type="checkbox"/> have not been made and will not be made.</li> </ol> </li> <li><input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3))</li> <li><input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4))</li> <li><input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).</li> </ol> <p><b>Items 11. to 16. below concern document(s) or information included:</b></p> <ol style="list-style-type: none"> <li><input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98</li> <li><input type="checkbox"/> An Assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.</li> <li><input type="checkbox"/> A <b>FIRST</b> preliminary amendment. <input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment.</li> <li><input type="checkbox"/> A substitute specification</li> <li><input type="checkbox"/> A change of power of attorney and/or address letter</li> <li><input checked="" type="checkbox"/> Other items or information             <ul style="list-style-type: none"> <li><input checked="" type="checkbox"/> Courtesy copy of the first page of the International Publication (WO 01/13951)</li> <li><input checked="" type="checkbox"/> Courtesy copy of the International Preliminary Examination Report (In Japanese).</li> <li><input checked="" type="checkbox"/> Formal drawings, 6 sheets, Figures 1-7</li> <li><input checked="" type="checkbox"/> Courtesy Copy of the International Search Report</li> </ul> </li> </ol> <p><input checked="" type="checkbox"/> The application is (or will be) assigned to <b>CHUGAI SEIYAKU KABUSHIKI KAISHA</b>, whose address is 5-1, Ukima 5-chome, Kita-ku, Tokyo 115-8543, Japan</p>		

JC13 Rec'd PCT/PTO 19 FEB 2002

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)		International Application No		Attorney's Docket No																																													
10/049957		PCT/JP00/05590		KATO=21																																													
<p>17. [xx] The following fees are submitted</p> <p><b>BASIC NATIONAL FEE (37 CFR 1.492 (a)(1)-(5):</b></p> <p>Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO. .... ..\$1040.00</p> <p>International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO .. \$890.00</p> <p>International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO. .... \$740.00</p> <p>International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) ....\$710.00</p> <p>International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4)... ..\$100.00</p> <p><b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b></p> <p>Surcharge of \$130.00 for furnishing the oath or declaration later than [ ] 20 [ ] 30 months from the earliest claimed priority date (37 CFR 1.492(e)).</p> <table border="1"><thead><tr><th>Claims as Originally Presented</th><th>Number Filed</th><th>Number Extra</th><th>Rate</th></tr></thead><tbody><tr><td>Total Claims</td><td>24 - 20</td><td>04</td><td>X \$18 00</td></tr><tr><td>Independent Claims</td><td>6 - 3</td><td>03</td><td>X \$84 00</td></tr><tr><td colspan="3">Multiple Dependent Claims (if applicable)</td><td>+ \$280.00</td></tr><tr><td colspan="3"><b>TOTAL OF ABOVE CALCULATIONS =</b></td><td>\$1,494.00</td></tr></tbody></table> <table border="1"><thead><tr><th>Claims After Post Filing Prel Amend</th><th>Number Filed</th><th>Number Extra</th><th>Rate</th></tr></thead><tbody><tr><td>Total Claims</td><td>- 20</td><td></td><td>X \$18.00</td></tr><tr><td>Independent Claims</td><td>- 3</td><td></td><td>X \$84 00</td></tr><tr><td colspan="3"><b>TOTAL OF ABOVE CALCULATIONS =</b></td><td>\$1,494.00</td></tr></tbody></table> <p>Reduction of 1/2 for filing by small entity, if applicable. Applicant claims small entity status. See 37 CFR 1.27</p> <p><b>SUBTOTAL =</b> \$1,494.00</p> <p>Processing fee of \$130.00 for furnishing the English translation later than [ ] 20 [ ] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).</p> <p><b>TOTAL NATIONAL FEE =</b> \$1,494 00</p> <p>Fee for recording the enclosed assignment (37 CFR 1.21(h)) The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) \$40.00 per property +</p> <p><b>TOTAL FEES ENCLOSED =</b> \$1,494.00</p> <table border="1"><tr><td rowspan="2"></td><td>Amount to be:</td><td>\$</td></tr><tr><td>refunded</td><td></td></tr><tr><td></td><td>charged</td><td>\$</td></tr></table>				Claims as Originally Presented	Number Filed	Number Extra	Rate	Total Claims	24 - 20	04	X \$18 00	Independent Claims	6 - 3	03	X \$84 00	Multiple Dependent Claims (if applicable)			+ \$280.00	<b>TOTAL OF ABOVE CALCULATIONS =</b>			\$1,494.00	Claims After Post Filing Prel Amend	Number Filed	Number Extra	Rate	Total Claims	- 20		X \$18.00	Independent Claims	- 3		X \$84 00	<b>TOTAL OF ABOVE CALCULATIONS =</b>			\$1,494.00		Amount to be:	\$	refunded			charged	\$	<p><b>CALCULATIONS</b> PTO USE ONLY</p>	
				Claims as Originally Presented	Number Filed	Number Extra	Rate																																										
				Total Claims	24 - 20	04	X \$18 00																																										
				Independent Claims	6 - 3	03	X \$84 00																																										
				Multiple Dependent Claims (if applicable)			+ \$280.00																																										
				<b>TOTAL OF ABOVE CALCULATIONS =</b>			\$1,494.00																																										
				Claims After Post Filing Prel Amend	Number Filed	Number Extra	Rate																																										
				Total Claims	- 20		X \$18.00																																										
				Independent Claims	- 3		X \$84 00																																										
				<b>TOTAL OF ABOVE CALCULATIONS =</b>			\$1,494.00																																										
	Amount to be:	\$																																															
	refunded																																																
	charged	\$																																															
<p>a. [ ] A check in the amount of \$ to cover the above fees is enclosed</p> <p>b. [X] Credit Card Payment Form (PTO-2038), authorizing payment in the amount of \$ 1,494 00, is attached.</p> <p>c. [ ] Please charge my Deposit Account No. 02-4035 in the amount of \$ to cover the above fees.</p> <p>A duplicate copy of this sheet is enclosed</p> <p>d. [XX] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 02-4035 A duplicate copy of this sheet is enclosed.</p> <p><b>NOTE:</b> Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.437(a) or (b)) must be filed and granted to restore the application to pending status.</p> <p>SEND ALL CORRESPONDENCE TO.</p> <p><b>BROWDY AND NEIMARK, P.L.L.C.</b> 624 NINTH STREET, N.W., SUITE 300 WASHINGTON, D.C. 20001 TEL: (202) 628-5197 FAX: (202) 737-3528 Date of this submission: February 19, 2002</p>				<p> SIGNATURE Sheridan Neimark NAME 20,520 REGISTRATION NUMBER</p>																																													

6/p/s

## SPECIFICATION

CHONDROGENESIS STIMULATORTECHNICAL FIELD

This invention relates to a novel chondrogenesis  
5 stimulator. More specifically, the invention relates to a  
chondrogenesis stimulator containing a membrane-bound  
transferrin-like protein (hereunder sometimes referred to  
as MTf).

PRIOR ART

10 The cartilage tissue of animals is composed of  
chondrocytes and matrix. The cartilage tissue accounts  
for the greater part of the skeleton at the prenatal stage  
and it is postnatally replaced with bone tissue due to  
endochondral ossification. When endochondral ossification  
15 starts, chondrocytes change from the resting to  
proliferating phase and the proliferating chondrocytes  
are then differentiated into hypertrophic chondrocytes  
[Reference; , "Hone no kagaku (Science of Bone)", ed. by  
Tsuda et al., pp. 11-29, Tokyo Ishiyaku Shuppan, 1982].  
20 Thus, it has been well known that chondrocytes are  
essential cells for the formation of bone tissue,  
particularly at the growth stage. However, the  
differentiation of chondrocytes and the endochondral  
ossification remain unknown in many aspects.

25 The cell membrane of chondrocytes has characteristic  
glycoproteins and their membrane proteins might contribute  
to the unique features of chondrocytes that distinguish  
them from the cells of other connective tissues (as

exemplified by spherical cell morphology, massive secretion of cartilage matrix, survival and proliferation in soft agar). Based on this hypothesis, Yan et al. (Yan et al.; J. Biol. Chem., vol. 265, pp. 10125-10131, 1990) and Kato et al. (Kato et al., Journal of the Society of Bone Metabolism of Japan, vol. 10, No. 2, pp. 187-192, 1992) investigated the effects of various lectins on the differentiation and proliferation of chondrocytes and, among other things, they have shown that concanavalin A (hereunder sometimes referred to as Con A) which is Jack bean lectin and which has affinity for  $\alpha$ -D-mannose residue and  $\alpha$ -D-glucose residue is a potent stimulator of chondrogenic differentiation, with the increase in proteoglycan synthesis being one of the criteria for the Con A activity. Chondrocytes treated with Con A change their shape from the immature flat morphology to the differentiated spherical form, inducing the production of proteoglycan and type II collagen which are markers of chondrogenic differentiation, the expression of alkaline phosphatase, etc., and even the calcification. Other lectins do not exert such differentiation inducing action.

In an attempt to search for a receptor mediating the action of Con A, Kawamoto et al. (Kawamoto et al., Eur. J. Biochem. vol. 256, pp. 503-509, 1998) paid particular attention to a protein of 76 kDa (p76) which was one of the about 20 kinds of Con A-binding proteins on chondrocytes and which would be expressed at lower levels in retinoic acid treated chondrocytes (upon treatment with retinoic

acid, chondrocytes are dedifferentiated to lose reactivity with Con A). After purifying p76 from the plasma membrane fraction of rabbit chondrocytes by Con A affinity column chromatography, the N-terminal amino acid sequence was  
5 determined and the gene was cloned. In view of the determined amino acid sequence and the nucleic acid sequence of its cDNA, p76 showed 86% amino acid identity with melanotransferrin (p97) and was considered its counterpart; p97 is known as a tumor-associated antigen  
10 expressed at high levels in human tumors such as melanoma. The physiological functions of p97 are yet to be known and its expression has been reported to be high in only tumor cells, with very low detectability in normal tissue.

In view of its ability to bind with Con A, p76 is  
15 presumably involved in the differentiation of chondrocytes or in the development of their function; however, nothing has been confirmed about the effects this protein would actually impose on chondrocytes or their precursors.

An object, therefore, of the present invention is to  
20 identify a substance that will be involved in the differentiation of chondrocytes and provide a novel chondrogenesis stimulator using the substance. The present invention will lead to the invention of a substance that can control the function of chondrocytes and which  
25 eventually enables promoted osteogenesis. The substance can potentially lead to the treatment, prevention and diagnosis of new types of diseases associated with the cartilage and bone metabolisms.

# DISCLOSURE OF THE INVENTION

In order to attain the stated object, the present inventors made intensive studies and found that differentiation to cartilage could be markedly induced by overexpressing a membrane-bound transferrin-like protein (MTf) gene in mouse cell line ATDC5 which retained the ability to differentiate to chondrocytes but which would hardly differentiate in the absence of stimulation.

Thus, the present invention provides a chondrogenesis stimulator containing a membrane-bound transferrin-like protein (MTf).

The MTf is preferably rabbit p76 protein, human p97 protein, mouse MTf protein, as well as a protein demonstrating the MTf activity that has an amino acid sequence encoded by DNA which hybridizes, under stringent conditions, with a DNA that encodes p76 protein or p97 protein or mouse MTf, and human p97 protein is particularly preferred.

The MTf is most preferably selected from the following:

- 1) a protein having the amino acid sequence of SEQ ID NO: 2;
- 2) a protein having the amino acid sequence of SEQ ID NO: 4;
- 3) a protein having the amino acid sequence of SEQ ID NO: 15; and
- 4) a protein demonstrating the MTf activity that has an amino acid sequence encoded by DNA which hybridizes, under

stringent conditions, with a DNA encoding the protein of  
SEQ ID NO: 2, 4 or 15.

The present invention also provides said  
chondrogenesis stimulator in which the MTf lacks the GPI  
5 anchor region.

The chondrogenesis stimulator of the invention  
becomes more effective when used in combination with an MTf  
activating agent and/or insulin.

The chondrogenesis stimulator of the invention is  
10 useful with the following diseases: OA (osteoarthritis);  
RA (rheumatoid arthritis); injury of articular cartilage  
due to trauma; maintenance of chondrocyte phenotype in  
autologous transplantation of chondrocytes; reconstruction  
of cartilage in the ear, trachea or nose; osteochondritis  
15 dissecans; regeneration of intervertebral disk or meniscus;  
fractured bone; and osteogenesis from cartilage.

The invention further provides an agent for gene  
therapy to promote chondrogenesis which contains as an  
active ingredient an expression vector incorporating a DNA  
20 coding for any one of the following proteins:

- 1) a protein having the amino acid sequence of SEQ ID NO:  
2;
- 2) a protein having the amino acid sequence of SEQ ID NO:  
4;
- 25 3) a protein having the amino acid sequence of SEQ ID NO:  
15; 4) a protein demonstrating the MTf activity that has an  
amino acid sequence encoded by DNA which hybridizes, under  
stringent conditions, with a DNA encoding the protein of

SEQ ID NO: 2, 4 or 15; and

5) a protein which is the same as protein 1), 2), 3) or 4), except that it lacks the GPI anchor region.

The present invention further provides a chondrogenic  
5 differentiation suppressing agent containing an MTf antagonist.

The MTf antagonist is preferably an anti-MTf antibody or an oligonucleotide or an oligonucleotide analog that are hybridizable with a nucleic acid encoding MTf.

10 The present invention further provides a method for screening an MTf activating agent which comprises the steps of:

1) preparing a cell line in which MTf is overexpressed, wherein said cell line retains the ability to differentiate  
15 to chondrocytes but hardly differentiate without stimulation;

2) adding candidate substances to the cell line prepared in step 1) and culturing it for a specified period of time; and

20 3) examining the cell line for induced chondrogenic differentiation and selecting an MTf activating agent from the candidate substances.

The present invention also provides an MTf activating agent as obtained by the method described above.

25 The present invention also provides a chondrogenesis stimulator containing an MTf activating agent as obtained by the method described above.

The present invention further provides MTf which



lacks the GPI anchor region.

# BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows a scheme of the procedure of preparing an MTf overexpressing ATDC5 variant cells;

5 Fig. 2 shows the expression of the MTf gene in the variant of ATDC5 cells as analyzed by Northern blotting (photographs of electrophoresis);

Fig. 3 shows the expression of the MTf protein in the variant of ATDC5 cells as analyzed by Western blotting  
10 (photographs of electrophoresis);

Fig. 4 is a set of photographs showing that MTf overexpressing cell lines (Full-1 and Full-5) demonstrate the morphology of differentiated chondrocytes in comparison with control cells (pC-1), all of which are cultured for 29  
15 days in the absence of insulin (biological morphology is shown);

Fig. 5 is a set of photographs showing that MTf overexpressing cell lines (Full-1 and Full-5) demonstrate the morphology of differentiated chondrocytes in comparison  
20 with control cells (pC-1), all of which are cultured for 29 days in the presence of insulin (biological morphology is shown);

Fig. 6 is a set of photographs showing the effects of the addition of the conditioned medium of rabbit  
25 chondrocytes on the induction of chondrogenic differentiation (biological morphology is shown); and

Fig. 7 shows the result of RT-PCR Southern blotting which demonstrates the overexpression of antisense MTf RNA

and the suppression of aggrecan synthesis in the presence and absence of insulin.

BEST MODE FOR CARRYING OUT THE INVENTION

The term "membrane-bound transferrin-like protein  
5 (MTf)" as used in the invention means a protein on the cell membrane of chondrocytes that binds to Con A and which has iron-binding sites as transferrin does. Preferably, the term means a protein having the ability to mediate the induction of chondrogenic differentiation by Con A.

10 The term MTf has conventionally been used as the abbreviation for melanotransferrin (p97) known as a tumor antigen expressed at high levels in melanoma and other tumors. As it turned out, however, p97 is also expressed at high levels in tissues other than cancer, particularly  
15 in cartilage. Since p97 is by no means specific to cancer tissue, the present inventors redefined the term MTf as meaning "membrane-bound transferrin-like protein".

The term "MTf activity" as used herein means an activity that induces undifferentiated cells to  
20 differentiate to cartilage and which promotes chondrocytes to develop their function.

Examples of MTf include but are not limited to rabbit p76 protein, p97 protein which is a human protein homologous to rabbit p76 protein, mouse MTf protein, as  
25 well as proteins having MTf activity that contain alterations such as deletion, substitution or addition of one or more of the amino acids of these proteins, and proteins having MTf activity amino acid sequences encoded

by DNA which hybridizes with DNA encoding p76 protein or  
p97 protein or mouse MTF protein under stringent conditions  
[a typical method is described in Molecular Cloning: A  
Laboratory Manual, Sambrook et al., Cold Spring Harbor  
5 Laboratory Press, 1989 and consists, for example, of  
hybridization at 68°C in a solution containing 6 x SSC,  
0.5% SDS, 10 mM EDTA, 5 x Denhardt's solution and 10 mg/ml  
of denatured salmon sperm DNA].

Rabbit p76 protein is homologous to human p97 protein  
10 and sometimes called rabbit p97 (Kawamoto et al., Eur. J.  
Biochem. vol. 256, pp. 503-509, 1998). The nucleotide and  
amino acid sequences of rabbit p76 protein are identified  
by SEQ ID NO: 1 and SEQ ID NO: 2, respectively. The  
nucleotide and amino acid sequences of human p97 protein  
15 are also known (Rose, T.M. et al., Proc NAS 83, 1261-1265,  
1986). The nucleotide and amino acid sequences of human  
p97 protein are identified by SEQ ID NO: 3 and SEQ ID NO:  
4, respectively. Mouse MTF protein is described in  
Biochim. Biophys. Acta, 1447:258-264, 1999 and its  
20 nucleotide and amino sequences are identified by SEQ ID NO:  
14 and SEQ ID NO: 15, respectively. The homology between  
the MTF proteins over animal species are high and the amino  
acid identity is 83% between mouse and human, 82% between  
mouse and rabbit, and 86% between human and rabbit.

25 p76/p97 proteins are GPI anchored proteins which have  
glycolipid GPI (glycosylphosphatidylinositol) bound to the  
carboxyl group in C-terminal amino acid so that they are  
bound to membranes using GPI as an anchor (for p76, see Ryo

Oda, Journal of Dentistry, Hiroshima University, vol. 29, No. 1, pp. 40-57, 1997; for p97, see Alemany, R. et al., J. Cell Science, 104, 1155-1162, 1993). As will be shown later in the Examples, it was verified that not only  
5 full-length MTf but also GPI anchor lacking MTf or soluble MTf have a chondrogenic differentiation inducing activity when they were expressed in non-MTf-expressing cells. Therefore, such soluble MTf, preferably the GPI anchor lacking MTf, can also be used as a chondrogenesis  
10 regulating agent. The GPI anchor lacking MTf as used herein means a soluble MTf which lacks all or part of the GPI anchor moiety; in the case of rabbit MTf, it may be exemplified by MTf in which the 28 residues at C-terminal necessary for GPI anchor binding are deleted and in the  
15 case of human MTf and mouse MTf, it may be exemplified by MTf in which the 30 residues at C-terminal necessary for GPI anchor binding are deleted.

The MTf to be used in the invention may be of a native or recombinant form and either form can be obtained  
20 by methods known in the art. The respective types of MTf are illustrated below.

#### Native form

MTf can be prepared by the method described in JP 7-82297A using chondrocytes. Briefly, cartilage tissues of  
25 various animals can be used as chondrocyte source; for example, a rabbit costal growth plate cartilage as the source is treated with protease and collagenase in accordance with the method of Kato et al. (Kato et al.; J.

Cell Biol., vol. 100, pp. 477-485, 1985) to obtain chondrocytes. The isolated chondrocytes can be incubated in a medium containing fetal calf serum (FCS) on a culture dish at 37°C in an atmosphere of 5% CO<sub>2</sub> and 95% air. The  
5 cultured chondrocytes are recovered, homogenized with a homogenizer and subjected to sedimentation equilibrium centrifugation by 17%/40% sucrose equilibrium density gradient to separate membrane proteins. The obtained membrane protein fraction is directly subjected on a  
10 concanavalin A affinity column; alternatively, in order to remove membrane proteins that bind to lectins other than concanavalin A, the membrane protein fraction is first subjected on an affinity column of wheat germ lectin which is a typical lectin and then subjected on a concanavalin A  
15 affinity column. By these and other techniques, more of the concanavalin A binding protein fraction can be separated. The specificity of the obtained concanavalin A bound protein fractions for chondrocytes can be evaluated by comparing these fractions through SDS-polyacrylamide gel  
20 electrophoresis (SDS-PAGE). After identifying the desired chondrocyte specific glycoproteins, bands of interest are excised from the gel, extracted and purified by electroelution or other suitable techniques. The resulting glycoproteins can be analyzed for the sugar chains after  
25 excising by endoglicosidase.

#### Recombinant form

Recombinant MTf can be prepared by the methods described in the Examples of the invention or modifications

thereof; by these methods, plasmids incorporating the MTf gene are transfected to host cells for expressing the MTf protein.

5        However, these are not the only methods that can be used and various methods of transformation and various host cells that are known in the art can also be used. For example, a gene encoding MTf may be inserted into a suitable vector to transform prokaryotic or eukaryotic host cells.

10       Further, suitable stimulators or sequences that are involved gene expression may be introduced into the vectors to enable gene expression in the transformed host cells. If desired, a gene of interest may be linked to genes encoding other polypeptides to express it as a fused  
15       protein so that it can be purified with greater ease or expressed at higher level. The desired protein can also be excised by applying suitable treatments in the purification step.

20       It is generally held that eukaryotic genes show polymorphism as is known for the human interferon gene. The polymorphism may cause substitution by one or more amino acids or it may cause changes in base sequences with no change occurring in amino acids.

25       Even polypeptides having deletion or addition of one or more amino acids within the amino acid sequence of SEQ ID NO: 2, 4 or 15 or having substitution of one or more amino acids may have a cell cycle regulating activity. For example, it is already known that a polypeptide having

substitution of cysteine for serine in the human interleukin 2 (IL-2) which is derived from nucleotide alterations has exerted an IL-2 activity (Wang et al., Science 224:1431, 1984). These techniques for preparing  
5 modified genes encoding MTf protein are known to the skilled artisan.

In many cases of expression in eukaryotic cells, sugar chains may be added to the protein and the addition of sugar chains can be regulated by substituting one or  
10 more amino acids of the protein and even in this case, the chondrogenic differentiation inducing activity may be exhibited. Therefore, genes encoding such polypeptides obtained by using artificial modifications of the gene encoding MTf gene can all be used in the invention, as long  
15 as such polypeptides have the chondrogenic differentiation inducing activity.

Expression vectors that can be used include replication origins, selection markers, promoters, RNA splicing sites, polyadenylation signals and so on.

20 Prokaryotic organisms that can be used as host cells in the expressing system include, for example, *Escherichia coli* and *Bacillus subtilis*. Eukaryotic microorganisms that can be used as host cells include, for example, yeasts and myxomycetes. If desired, insect cells such as Sf9 may be  
25 used as host cells. Host cells derived from animal cells include, for example, COS cells and CHO cells.

Transformants thus obtained by transforming with the gene encoding MTf protein are cultured to produce proteins.

The proteins can be recovered from the transformants or from the cultured medium and be purified. Not only the proteins which are obtained with genes containing nucleotide sequences encoding the amino acid sequences of SEQ ID NO: 2, 4 and 15 but also proteins which are obtained using genes containing altered nucleotide sequences encoding the amino acid sequences having substitution, deletion or addition of one or more amino acids within the amino acid sequences of SEQ ID NO: 2, 4 and 15, or proteins which are obtained using nucleotide sequences that hybridize those altered nucleotide sequences can be used as the chondrogenesis promoter of the invention as long as they have the biological function of MTf protein, namely, the chondrogenic differentiation inducing activity.

Conventional methods for separating and purifying proteins can be employed to separate and purify the MTf protein. For example, various techniques of chromatography, ultrafiltration, salting-out, dialysis, etc. can appropriately be selected and used in combination.

To use the chondrogenesis stimulator of the invention, the MTf described above may be administered in the form of a protein or it may be used an agent for gene therapy.

Insulin or an insulin-like growth factor has conventionally been known as a chondrocyte differentiating substance. It is interesting to note that the chondrogenesis stimulator of the invention induces chondrogenic differentiation even in the absence of



insulin. However, it was found that the effect of the chondrogenesis stimulator is further enhanced in the presence of insulin. Therefore, the desired cartilage repairing action could be further enhanced by using MTF in  
5 combination with MTF activating agents such as insulin and an insulin-like growth factor.

When the supernatant of a chondrocyte culture was added, marked differentiation of chondrocytes was observed in MTF overexpressing cell lines, suggesting that an MTF  
10 activating agent may exist in the conditioned medium of a chondrocyte culture. Therefore, the desired cartilage repairing action could be further potentiated by using MTF in combination with an MTF activating agent.

The MTF activating agent may be obtained by the  
15 following methods:

- 1) purifying from the conditioned medium of a chondrocyte culture system;
- 2) cloning the cDNA for protein binding to an MTF from a chondrocyte cDNA library; and
- 20 3) cloning the cDNA of protein binding to an MTF by the yeast two-hybrid method.

To screen various candidate substances for an MTF activating agent, a method including the following steps can be used:

- 25 1) preparing a cell line in which MTF is overexpressed, wherein said cell line retains the ability to differentiate to chondrocytes but hardly differentiate without stimulation;



MTf protein or an MTf mutant (variant) introduced in it. The agent for gene therapy of the invention contains as an active ingredient an expression vector incorporating a DNA coding for any one of the following proteins:

- 5 1) a protein having the amino acid sequence of SEQ ID NO: 2;
- 2) a protein having the amino acid sequence of SEQ ID NO: 4;
- 3) a protein having the amino acid sequence of SEQ ID NO: 15;
- 10 4) a protein demonstrating the MTf activity that has an amino acid sequence encoded by DNA which hybridizes, under stringent conditions, with a DNA encoding the protein of SEQ ID NO: 2, 4 or 15; and
- 5) a protein which is the same as protein 1), 2), 3) or 4),  
15 except that it lacks the GPI anchor region.

DNAs encoding MTf variants can be easily prepared by the skilled artisan using known techniques such as site-directed mutagenesis and PCR [Molecular Cloning: A Laboratory Manual, 2nd ed., Chapter 15, Cold Spring Harbor Laboratory Press (1989), and PCR - A Practical Approach, 20 IRL Press, 200-210 (1991)].

In the present invention, an MTf- or MTf variant expression vector is provided as a DNA to be introduced into cells. These expression vectors can be prepared by  
25 linking DNA encoding an MTf- or MTf variant to an expression vector such as pSG5 (Stratagene). In the next step, the prepared DNA mixture is introduced into cells. Exemplary cells may include bone marrow interstitial cells,

fibroblasts, periosteal cells, perichondral cells, synovial cells and dedifferentiated chondrocytes. DNA can be introduced into cells by the calcium phosphate method [Idenshi donyu to hatsugen kaisekiho (Gene Introduction and Methods of Expression and Analysis), ed. by Takashi Yokota and Kenichi Arai, Yodosha, 1994]. Hence, by using the introduced DNA as a medicinal active ingredient, one can prepare an agent for gene therapy which has the chondrogenesis promoting action. It is thought that, upon administering such agent for gene therapy, MTF or its variant would be expressed at high levels in cells, promoting the action of inducing chondrogenic differentiation in the cells. Therefore, the MTF containing agent of the invention for gene therapy can be used as a therapeutic or preventive of the various diseases listed above.

The agent of the invention for gene therapy can be introduced into cells by either a virus vector based method of gene introduction or a non-viral method of gene introduction [Nikkei Science, April 1994, pp. 20-45, Jikken igaku zokan (Extra Issue of Experimental Medicine), 12(15)(1994), and Jikken igaku bessatsu (Supplement to Experimental Medicine), "Idenshi chiryo no kiso gijutsu (Basic Technology in Gene Therapy)", Yodosha (1996)].

In an example of the viral vector based method of gene introduction, DNA encoding an MTF or a variant MTF is inserted into DNA or RNA viruses such as retrovirus, adenovirus, adeno-associated virus, herpes virus, vaccinia

virus, poxvirus, poliovirus and Sindbis virus. Non-viral methods of gene introduction include direct intramuscular administration of an expression plasmid (DNA vaccination), liposome method, lipofectin method, microinjection, calcium phosphate method, and electroporation.

In order to ensure that the agent for gene therapy of the invention acts as a practical medicine, two methods may be used, that is, an in vivo method where DNA is directly introduced into the body and an ex vivo method where a certain kind of cells are taken out of a human and DNA is introduced into the cell, which is then put back into the body [Nikkei Science, April 1994, pp. 20-45, Gekkan yakuji (Monthly Yakuji), 36(1), 23-48 (1994), and Jikken igaku zokan (Extra Issue of Experimental Medicine), 12(15)(1994)]. The in vivo method is more preferred.

When administering the agent for gene therapy by the in vivo method, the route of administration should depend on the disease, its severity and other factors. Exemplary methods of administration include intra-articular injection, direct application to a missing part of articular cartilage, implantation (with putty, polylactic acid, etc.) and intra-articular sustained-release agent. Intravenous injection is also possible. For administration by the in vivo method, injections are generally used, with conventional carriers being added as required. Liposomes or membrane fused liposomes may be formulated as suspensions, frozen vesicles, centrifugally concentrated frozen vesicles.



amino acid sequence identified by SEQ ID NO: 2, 4 or 15.

The term "oligonucleotide" as used herein means oligonucleotides generated from naturally occurring bases and sugar portions bound by intrinsic phosphodiester bonds, as well as analogs thereof. Therefore, the first group encompassed by this term comprises naturally occurring species or synthetic species that are generated from naturally occurring subunits or homologs thereof. The term "subunit" means a base-sugar combination which links to adjacent subunit by phosphodiester bond or other bond. The second group of oligonucleotides are their analogs that function similar to oligonucleotides but which are composed of residues having non-naturally-occurring moieties. These include oligonucleotides having chemical modifications applied to phosphate groups, sugar portions and 3'- and 5'-ends in order to provide increased stability. Examples are oligophosphorothioate and oligomethylphosphonate in which one of the oxygen atoms in the phosphodiester group between nucleotides is substituted by sulfur and  $-CH_3$ , respectively. Phosphodiester bonds may be replaced by other structures which are non-ionic and achiral. Additional oligonucleotide analogs that can be used are species containing modified base forms, that is, purine and pyrimidine in non-naturally-occurring form.

The oligonucleotides to be used in the invention have preferably 5 - 40 subunits in length, more preferably 8 - 30 subunits, most preferably 12 - 30 subunits.

In the present invention, the target portion of mRNA

with which oligonucleotides hybridize is preferably a transcription initiation site, a translation initiation site, an intron/exon junction site or a 5'-cap site; considering the secondary structure of mRNA, a site having  
5 no steric hindrance should be selected.

In the present invention, peptide nucleic acids (see, for example, Bioconjugate Chem., Vol. 5, No. 1, 1994) may be used in place of oligonucleotides.

In a particularly preferred embodiment of the  
10 invention, oligonucleotides or peptide nucleic acids that hybridize with a nucleotide sequence encoding the amino acid sequence identified by SEQ ID NO: 2 and which can inhibit MTF expression is employed.

In the present invention, oligonucleotides can be  
15 produced by synthesis methods known in the art, for example, the solid-phase synthesis method using a synthesizer as manufactured by Applied Biosystems. Similar methods can be used to produce oligonucleotide analogs such as phosphorothioate and alkylated derivatives [Akira  
20 Murakami et al., "Kinosei antisense DNA no gosei (Chemical Synthesis of Functional Antisense DNA)", Organic Synthesis Chemistry, 48(3):180-193, 1990].

The MTF antagonist that can be used in the invention is not limited to oligonucleotides of the above-defined  
25 antisense DNA providing length. To the extent that production of intrinsic MTF can be suppressed, a longer antisense, preferably an antisense of 500 - 600 nucleotides in length, may be inserted into a genome to be used for



suppressing chondrogenic differentiation (see Example 3).

The anti-MTf antibody to be used in the invention is one that recognizes a peptide having at least five consecutive amino acids in the amino acid sequence identified by SEQ ID NO: 2, 4 or 15; this can be produced using a conventional procedure [see, for example, Shin-seikagaku jikken koza 1 (New Course in Biochemical Experiments 1), Protein I, pp. 389-397, 1992], which comprises immunizing an animal with an antigenic peptide having at least five consecutive amino acids in the amino acid sequence of SEQ ID NO: 2, 4 or 15, isolating the antibody produced in the animal body, and purifying the isolated antibody. The antibody may include a polyclonal and a monoclonal antibody and methods of preparing these antibodies are also known to the skilled artisan.

The following examples are provided to further illustrate the present invention but are in no way to be taken as limiting the invention. Various alterations and modifications can be made by the skilled artisan and are included within the scope of the invention.

### Examples

#### Materials and Methods of Experiment

##### Rabbit chondrocyte culture

Chondrocytes were isolated from rabbit costal cartilage using, with necessary modifications, the method of Kato et al. (Kato et al.: J. Cell Biol., vol. 100, pp. 477-485, 1985). Specifically, the resting cartilage of ribs in 4-week old male Japanese albino rabbits (Hiroshima

Laboratory Animals) was separated, shredded with a surgical knife and incubated in a Dulbecco's modified Eagle's medium (DMEM, Flow Laboratories) containing 8 mg/mL of actinase E (Kaken Seiyaku) and 5% fetal calf serum for 1 hour and in  
5 DMEM containing 0.15% collagenase (Worthington Biochemical) for 3 hours. Cells passing through a 120- $\mu$ m nylon filter were recovered, seeded in plastic culture dishes (Corning) and grown in an alpha-modified Eagle's medium ( $\alpha$ -MEM, Sanko Junyaku) containing 10% fetal calf serum (Mitsubishi  
10 Kasei), 50  $\mu$ g/mL of ascorbic acid, 50 U/mL of G potassium, 60  $\mu$ g/mL of kanamycin (all being from Meiji Seika) and 250  $\mu$ g/mL of amphotericin B (ICN Biochemical) (medium A) or in a serum-free DMEM (medium B) at 37°C in the atmosphere of 5% CO<sub>2</sub> gas.

15 Mouse chondrogenic cell line ATDC5 culture

ATDC5 was purchased from Riken Cell Bank (Tsukuba, Japan). The cells were cultured in a 1:1 mixture of Ham F-12 medium (Flow Laboratories) and a Dulbecco's modified Eagle's medium (DMEM, Flow Laboratories) containing 5%  
20 fetal calf serum (FCS, Mitsubishi Kasei), 10  $\mu$ g/mL of human transferrin (Boehringer Mannheim) and 0.3 nmol/mL of sodium selenite (Wako Junyaku) (maintenance medium) at 37°C in the atmosphere of 5% CO<sub>2</sub> gas. To induce cartilage differentiation, ATDC5 cells were cultured in a medium  
25 (differentiation medium) prepared by adding 10  $\mu$ g/mL of bovine insulin (Sigma) to the maintenance medium.

cDNA Cloning of rabbit MTf and nucleotide sequencing

Three days after reaching confluence, the rabbit



### Creating MTf overexpressing ATDC5 variant cells

Rabbit MTf cDNA (Kawamoto T. et al., EJB, 1988) of either full length or a truncated form which had the 28 residues from C-terminal necessary for GPI anchor binding  
5 deleted was inserted into pcDNA3.1/Zeo(+) plasmid expression vector (containing a cytomegalovirus very early promoter/enhancer; Invitrogen, San Diego, CA). Specifically, an EcoRI-NotI fragment including the full length was excised from a vector and inserted at the EcoRI-  
10 NotI site of pcDNA3.1/Zeo(+). To create a variant which lacks the GPI anchor binding site, a fragment was prepared having a stop codon inserted 28 amino acids upstream of the C-terminal and after confirming its sequence, the fragment was inserted at the EcoRI-NotI site of pcDNA3.1/Zeo(+).

15 In these ways, there were prepared a plasmid having a full length MTf cDNA (MTf Full) as an insert and a plasmid having GPI anchor-lacking MTf cDNA (MTf(-)GPI) as an insert; the two plasmids (pMTf Full and pMTf(-)GPI) were each transfected to ATDC5 cells (Riken, Tsukuba, Japan)  
20 using SuperFect Transfection Reagent (QIAGEN). By selection with Zeocin (Invitrogen), stable transformants were prepared.

Specifically,  $2 \times 10^5$  ATDC5 cells were seeded in 10-cm culture dishes. On the next day, 2  $\mu$ g each of the  
25 plasmid DNAs to be introduced (pMTf Full and pMTf(-)GPI) and about 40  $\mu$ L of SuperFect Transfection Reagent in solution were individually dissolved in a serum-free medium and stored until use, when they were rapidly mixed together

and added to ATDC5 cells washed with a serum-free medium. After incubation at 37°C for 1 hour in the atmosphere of 5% CO<sub>2</sub> gas, a serum-supplemented medium was added and cultivation was conducted for an additional day. A control  
5 group was prepared by transfecting only the vector.

One day after the transfection, selection was started in a serum-supplemented medium containing 50 µg/mL of Zeocin and cell culture was continued for 2 weeks with medium change effected on every third day. As a result,  
10 there were obtained ATDC5 variant cell lines that would assure stable expression of MTf Full and MTf(-)GPI and these variant cell lines were subcultured in a serum-supplemented medium containing 50 µg/mL of Zeocin.

A scheme of the procedure of creating MTf overexpressing ATDC5 variant cells is shown in Fig. 1.  
15

Expression of rabbit MTf gene in ATDC5 variant cell lines

Expression of rabbit MTf gene in ATDC5 variant cell lines was confirmed by Northern blotting. Specifically, total RNA was prepared from the ATDC5 variant cell lines by  
20 the guanidine thiocyanate method; 10 µg of the total RNA was electrophoresed on a 1% agarose gel containing 2.2 mol/L of formaldehyde and transferred onto Hybond-N membrane (Amersham). The membrane was hybridized with a <sup>32</sup>P labeled 2.2 kb rabbit MTf cDNA probe at 42°C for 16 hours.  
25 After washing the membrane, a BioMax X-ray film (Kodak) was exposed to the membrane at -80°C to detect signals. The result is shown in Fig. 2.

In the MTf Full cell line, it was found that the

rabbit MTf gene have been expressed strongly in clone Nos. 1, 4 and 5.

In the MTf(-)GPI cell line, it was found that the rabbit MTf gene have been expressed strongly in clone Nos. 3, 3N, 8, 9 and 10. In the Examples, (-)GPI-3 was used.

Expression of rabbit MTf protein in ATDC5 variant cell lines

Expression of rabbit MTf protein in ATDC5 variant cell lines was confirmed by Western blotting. Specifically, membrane fraction protein was prepared from the ATDC5 variant cell lines, subjected to SDS-PAGE at 10 µg/lane, and transferred to a polyvinylidene difluoride membrane (Milipore). After the transfer, the membrane was blocked with 4% skimmed milk and reacted with anti-MTf serum [1:500 dilution; Eur. J. Biochem, 256, 503-509 (1988)] at 4°C for 14 hours, then reacted with <sup>125</sup>I sheep anti-mouse IgG(Fab')<sub>2</sub> fragment (Amersham) at room temperature for 2 hours. The membrane was washed and a BioMax X-ray film was exposed to the membrane at -80°C for analysis. The result is shown in Fig. 3.

In the MTf Full cell line, it was found that the rabbit MTf protein have been expressed strongly in clone Nos. 1 and 5. These clones were named MTf overexpressing cell lines (Full-1 and Full-5).

Example 1: Chondrogenic differentiation in MTf overexpressing cell lines

The MTf overexpressed cell lines ( $4.0 \times 10^4$  cells) were seeded in 6-multiwell plates and cultured in a

maintenance medium at 37°C in a 5% CO<sub>2</sub> gas phase.

The MTf overexpressing cell lines to be investigated were Full-1 and Full-5 which were found to have expressed both the MTf gene and protein. The MTf(-)GPI cell line to  
5 be investigated was GPI-3 which was found to have expressed the MTf gene.

As control cells, ATDC5 cells and pC-1 (vector alone) were prepared in the same manner as described above and their morphological features were examined under a  
10 microscope. Cell morphology was examined with an Olympus phase-contrast microscope. Two microscopic fields were taken for each culture system and at least 200 cells were counted to calculate the proportion of round cells.

In the absence of insulin, the control cells (pC-1)  
15 did not differentiate to chondrocytes; on the other hand, the MTf overexpressing cell lines (Full-1 and Full-5) and the MTf(-)GPI cell line [(-)GPI-3] started to differentiate within 20 days and 29 days after seeding, almost all regions of the cells had differentiated to chondrocytes  
20 (Fig. 4).

The similar test was conducted in the presence of insulin (10 µg/mL) (insulin was added at day 0). The result was the same as in the absence of insulin (Fig. 5), except that, in the presence of insulin, further  
25 differentiation was induced in the MTf overexpressing cell lines, namely, more cells "rounded" like chondrocytes than in the absence of insulin.

The above results show the effectiveness of MTf in

inducing chondrogenic differentiation, which was exhibited even in the absence of insulin.

Example 2: Effect of Adding the Conditioned Medium of Rabbit Chondrocyte Culture

5           Resting rabbit chondrocytes ( $1 \times 10^6$  cells) were seeded in 10-cm culture dishes and cultured in medium A (10 mL) at 37°C in the atmosphere of 5% CO<sub>2</sub> gas. Two days after confluence, medium A was replaced by serum-free medium B (5 mL). After 24 hours of cell culture, the conditioned  
10 medium (CM) was recovered and subjected to an experiment. After the recovery, CM was supplemented with fetal calf serum at a concentration of 5%.

          MTf overexpressing cell line (Full-5) and the control cell line (pC-1) were seeded at  $8.0 \times 10^4$  cells in 6-  
15 multiwell plates and cultured in a maintenance medium at 37°C in the atmosphere of 5% CO<sub>2</sub> gas. Three days after confluence (day 7), the previously recovered CM was added to give a concentration of 60% in the overall liquid culture; at the same time, 10 µg/mL of bovine insulin was  
20 added. Cultivation was continued for additional 48 hours at 37°C in a 5% CO<sub>2</sub> gas phase.

          Forty-eight hours after the addition of CM, almost all cells of the MTf overexpressing cell line to which CM was added [Full-5(+)-CM] differentiated to chondrocytes  
25 which synthesized active substrate and which resembled paving stones. The cells of the MTf expressing cell line to which no CM was added [Full-5(-)-CM] had almost the same morphology as the control cell lines in which only a vector



was expressed [pC-2(+)CM and (-)CM]. The cell lines in which only a vector was expressed had no visible induced differentiation to chondrocytes due to the addition of CM (Fig. 6).

5           These results show the presence of an MTf activating agent in CM.

Example 3: Chondrogenic differentiation due to  
Overexpression of Antisense MTf RNA

ATDC5 variant cell lines (A-01, A-05, A-08, A-09, A-  
10 11, A-12, A-23 and A-24) in which mouse antisense MTf RNA was overexpressed were prepared by the similar method used in preparing the ATDC5 variant cell lines in which MTf was forcibly expressed. The mouse antisense MTf RNA was prepared as follows: cDNA fragments of mouse MTf were  
15 amplified by PCR (using primers 5'-  
GGTGTGTTGAGGGGCGTGGACTCT-3' (SEQ ID NO: 9) and 5'-  
TCACCAACGGCTTTGAGCACATCAC-3' (SEQ ID NO: 10), inserted into pGEM-T Easy Vector (Promega), excised with ApaI-NotI and inserted into pCDNA3.1/Zeo(+) at ApaI-NotI site (i.e.,  
20 inserted in reverse direction). The sequence of the inserted portion is identified by SEQ ID NO: 11.

A control group was also prepared by transfecting only the vector (pC-1).

The expression of mouse MTf antisense was examined by  
25 Northern blotting using the above-mentioned ApaI-NotI fragment as a probe.

A criterion for the suppression of chondrogenic differentiation was the suppression of synthesis of a

cartilage proteoglycan, aggrecan, and a test was conducted by RT-PCR Southern blotting both in the presence of insulin (added in an amount of 10 µg/mL after day 4) and in its absence. Specifically, total RNA was extracted from the  
5 cells of each clone by the guanidine thiocyanate method; single-stranded cDNA was synthesized from the extracted total RNA (1 µg) using SUPERScript pre-amplification system kit (Life Technologies); using the cDNA as a template, PCR was performed on the aggrecan gene with a pair of primers  
10 5'-TGCTACTTCATCGACCC-3' (forward) (SEQ ID NO: 12) and 5'-AAAGACCTCCCCTCCATCT-3' (reverse) (SEQ ID NO: 13); the PCR reaction mixture was electrophoresed on a 1% agarose gel and transferred onto Hybond-N membrane (Amersham). The membrane was hybridized with a <sup>32</sup>P labeled antisense MTF  
15 probe and mouse aggrecan cDNA probe at 42°C for 16 hours. The membrane was washed with 0.2 x SSC which contains 0.5% SDA and a BioMax X-ray film (Kodak) was exposed to the membrane at -80°C to detect signals.

The result is shown in Fig. 7. Antisense MTF RNA was  
20 expressed most strongly in variant cell line A-12 (lane 6), followed by A-11 (lane 5) in strength. Correspondingly, expression of the aggrecan gene was suppressed most effectively in variant cell line A-12 (lane 6), followed by A-11 (lane 5) in effectiveness. While expression of the  
25 aggrecan gene was suppressed in the absence of insulin, it was more effectively suppressed in the presence of insulin.



incorporating a DNA encoding any one of the following proteins:

- 1) a protein having the amino acid sequence of SEQ ID NO: 2;
- 2) a protein having the amino acid sequence of SEQ ID NO: 4;
- 3) a protein having the amino acid sequence of SEQ ID NO: 15;
- 4) a protein demonstrating the MTF activity that has an amino acid sequence encoded by DNA which hybridizes, under stringent conditions, with a DNA coding for the protein of SEQ ID NO: 2, 4 or 15; and
- 5) a protein which is the same as protein 1), 2), 3) or 4), except that it lacks the GPI anchor region.

8. The chondrogenesis stimulator according to claim 1 which is used in combination with an MTF activating agent.

9. The chondrogenesis stimulator according to claim 1 which is used in combination with insulin or an insulin-like growth factor.

10. The chondrogenesis stimulator according to any one of claims 1 - 9 for treating at least one bone disease selected from the following diseases in which chondrogenic differentiation is involved: OA (osteoarthritis); RA (rheumatoid arthritis); injury of articular cartilage due to trauma; maintenance of chondrocyte phenotypes in autologous chondrocyte transplantation; reconstruction of cartilage in the ear, trachea or nose; osteochondritis dissecans; regeneration of intervertebral disk or meniscus; bone fracture; and osteogenesis from cartilage.

11. A chondrogenic differentiation suppressing agent containing an MTf antagonist.

12. The chondrogenic differentiation suppressing agent according to claim 11, wherein the MTf antagonist is an anti-MTf antibody or an oligonucleotide or an oligonucleotide analog that are hybridizable with a nucleic acid encoding MTf.

13. A method for screening an MTf activating agent which comprises the steps of:

1) preparing a cell line in which MTf is overexpressed, wherein said cell line retains the ability to differentiate to chondrocytes but hardly differentiate without stimulation;

2) adding candidate substances to the cell line prepared in step 1) and culturing it for a specified period of time; and

3) examining the cell line for induced chondrogenic differentiation and selecting an MTf activating agent from the candidate substances.

14. An MTf activating agent obtained by the method according to claim 13.

15. A chondrogenesis stimulator containing an MTf activating agent obtained by the method according to claim 13.


16. MTf which lacks the GPI anchor region.

## ABSTRACT

There are provided a chondrogenesis stimulator containing MTf, a chondrogenic differentiation suppressing agent containing an MTf antagonist, a screening method for  
5 obtaining an MTf activating agent, an MTf activating agent obtained by the screening method, a chondrogenesis stimulator containing an MTf activating agent as obtained by the screening method, and MTf which lacks the GPI anchor region.

## Fig. 1

Construction of expression vector (pc-DNA 3.1 (+) plasmid)



GPI anchor region

Constructed expression vectors

MTf Full



MTf (-) GPI




Stable transfection




Checking for the expression of MTf mRNA by Northern blotting

Checking for the expression of MTf protein by Western blotting

*Fig.2*




*Fig.3*







*Fig.4*





*Fig.5*



*Fig.6*

*Fig.7***Antisense**

1 2 3 4 5 6 7 8

**MTf mRNA****day 23 (-) Insulin****AggreCan****day 17 (+) insulin****AggreCan**

1 2 3 4 5 6 7 8 9

- 1 = A-01
- 2 = A-05
- 3 = A-08
- 4 = A-09
- 5 = A-11
- 6 = A-12
- 7 = A-23
- 8 = A-24
- 9 = pC-1

100499571.021902  
 10/049,957  
 Rec'd PCT/PTO 05 JUL 2002

# SEQUENCE LISTING

<110> KATO, Yukio  
 FUJIMOTO, Katsumi

<120> CHONDROGENESIS PROMOTERS

<130> KATO=21

<140> 10/049,957

<141> 2002-02-19

<150> PCT/JS00/05590

<151> 2000-08-21

<150> JP 232966/1999

<151> 1999-08-19

<160> 15

<170> PatentIn version 3.1

<210> 1

<211> 2388

<212> DNA

<213> Oryctolagus cuniculus

<400> 1

gccgcccgtc actcggtcgc actcggactc agaccagtc cgacccctg gactgcgcca	60
tgcggtgccg aagcgcggct atgtggatct tcctggccct gcgcaccgca ctgggcagcg	120
tggagggtgcg gtgggtgcacc gcgtccgagc ccgagcagca gaagtgcgag gacatgagcc	180
aggccttccg cgaagccggc ctccagcccc cctgctgtg cgtgcagggc acctcgcccg	240
accactgcgt ccagctcatc gcggcccacg aggccgacgc catcactctg gacggaggag	300
ccatttacga ggcggggaag gaacacggcc tgaagcccg ggtgggagaa gtgtatgacc	360
aagagggtggg cacctcctac tacgctgtgg ccgtgggtcaa gaggagctcc aacgtgacca	420
tcaacaccct gagaggcgtg aagtcctgcc acacgggcat caaccgcacg gtgggctgga	480
acgtgcctgt gggctacctg gtggacagcg gccgcctctc agtgatgggc tgtgacgtgc	540
tcaaagcggc cagcgagtac ttccggggga gctgcgtccc tggggcagga gagaccagat	600
actcggagtc cctctgtcgc ctctgccggg gcgacacctc cggggagggg gtgtgtgaca	660
agagccccct ggagcggtag tacgactaca gcggggcctt ccggtgcctg gcagaaggcg	720
caggggacgt ggcctttgtg aagcacagca ccgtgctgga gaacacggat gggagaacac	780
tgcctcctg gggccacatg ctgatgtcac gggactttga gctgctgtgc cgggacggca	840
gccggggcag cgtcaccgag tggcagcact gccacctggc ccgggtgccc gccacgccc	900

tgggtgggtccg ggccgacacc gacgcaggcc tcattcttccg gcttctcaat gagggccagc 960  
 ggctgttcag ccacgagggc agcagcttcc agatgttcag ctccgaggcc tacggccaga 1020  
 agaacctgct gttcaaagac tccacgctgg agctgggtgcc catcgccaca cagacctacg 1080  
 aggcctggct gggccccgag tacctgcacg ccatgaaggg tctgctctgt gaccccaacc 1140  
 ggctgcccc atacctgcgc tgggtgcgtgc tgtccacccc cgagatccag aagtgtggag 1200  
 acatggccgt ggccttcagc cggcagaggc tcaagccgga gatccagtgt gtctcggcgg 1260  
 agtcccccca gcaactgcag gagcagatcc aggctgggca catcgatgct gtgacctga 1320  
 acggggagga cattcacaca gcggggaaga cttatgggct gatcccggtc gcgggggagc 1380  
 tgtatgcgc ggacgacagg agtaactcgt acttcgtggg ggccgtgggtg aagcgagaca 1440  
 ggcctacgc cttcaccgtg gacgagctgc gcgggaagcg ctctgcccac cccggcttcg 1500  
 gcagcccggc cggctgggac gtcccgtgg gcgcctcat ccaactggggc tacatccggc 1560  
 ccaggaactg cgacgtcctc acagcgggtg gtcagttctt caacgccagc tgtgtgccgg 1620  
 tgaacaaccc caagaagtac cctcctcgc tgtgcgcact ctgcgtgggt gacgagcagg 1680  
 gccgcaaca gtgcactggc aacagccagg agcgggtacta tggcgacagt ggcgccttca 1740  
 ggtgcctggg ggaggggtgca ggggacgtgg ccttcgtcaa gcacacgacc atctttgaca 1800  
 acacaaatgg ccacaatccc gagccgtggg ctgccatct gaggagccag gactacgagc 1860  
 tgctgtgccc caacggcgcg cgagctgagg cgcaccagtt tgccgcctgc aacctggccc 1920  
 agattccgtc ccacgccgtc atgggtgcggc ccgacaccaa catcttcacc gtttacggac 1980  
 tgctggacaa ggcccaggac ctgtttggag acgaccacaa caagaacggg ttcaagatgt 2040  
 tcgactcctc cagctaccac ggccgagacc tgctcttcaa ggacgccacg gtgcgcgctg 2100  
 tgctgtggg cgagaggacc acctaccagg actggctggg gccggactac gtggcggtc 2160  
 tggaagggat gcagtcacag cgggtgctcag gggcagccgt cggcgcccc ggggcctcgc 2220  
 tgctgccgt gctgcccctg gctgcgggccc tctgctgtc ttgctctga gagcagcccc 2280  
 gggcagcctc ggccccggca ggggagcctg cgcggaagct tctgaacga gcccgcgccc 2340  
 tggctggatg tggttacctc ggcgagccgc ggggcccgcg tcccccg 2388

<210> 2  
 <211> 736  
 <212> PRT  
 <213> Oryctolagus cuniculus  
 <400> 2

Met Arg Cys Arg Ser Ala Ala Met Trp Ile Phe Leu Ala Leu Arg Thr  
 1 5 10 15  
 Ala Leu Gly Ser Val Glu Val Arg Trp Cys Thr Ala Ser Glu Pro Glu  
 20 25 30  
 Gln Gln Lys Cys Glu Asp Met Ser Gln Ala Phe Arg Glu Ala Gly Leu  
 35 40 45  
 Gln Pro Ala Leu Leu Cys Val Gln Gly Thr Ser Ala Asp His Cys Val  
 50 55 60  
 Gln Leu Ile Ala Ala His Glu Ala Asp Ala Ile Thr Leu Asp Gly Gly  
 65 70 75 80  
 Ala Ile Tyr Glu Ala Gly Lys Glu His Gly Leu Lys Pro Val Val Gly  
 85 90 95  
 Glu Val Tyr Asp Gln Glu Val Gly Thr Ser Tyr Tyr Ala Val Ala Val  
 100 105 110  
 Val Lys Arg Ser Ser Asn Val Thr Ile Asn Thr Leu Arg Gly Val Lys  
 115 120 125  
 Ser Cys His Thr Gly Ile Asn Arg Thr Val Gly Trp Asn Val Pro Val  
 130 135 140  
 Gly Tyr Leu Val Asp Ser Gly Arg Leu Ser Val Met Gly Cys Asp Val  
 145 150 155 160  
 Leu Lys Ala Val Ser Glu Tyr Phe Gly Gly Ser Cys Val Pro Gly Ala  
 165 170 175  
 Gly Glu Thr Arg Tyr Ser Glu Ser Leu Cys Arg Leu Cys Arg Gly Asp  
 180 185 190  
 Thr Ser Gly Glu Gly Val Cys Asp Lys Ser Pro Leu Glu Arg Tyr Tyr  
 195 200 205  
 Asp Tyr Ser Gly Ala Phe Arg Cys Leu Ala Glu Gly Ala Gly Asp Val  
 210 215 220  
 Ala Phe Val Lys His Ser Thr Val Leu Glu Asn Thr Asp Gly Arg Thr  
 225 230 235 240  
 Leu Pro Ser Trp Gly His Met Leu Met Ser Arg Asp Phe Glu Leu Leu  
 245 250 255  
 Cys Arg Asp Gly Ser Arg Ala Ser Val Thr Glu Trp Gln His Cys His  
 260 265 270  
 Leu Ala Arg Val Pro Ala His Ala Val Val Val Arg Ala Asp Thr Asp  
 275 280 285  
 Ala Gly Leu Ile Phe Arg Leu Leu Asn Glu Gly Gln Arg Leu Phe Ser  
 290 295 300  
 His Glu Gly Ser Ser Phe Gln Met Phe Ser Ser Glu Ala Tyr Gly Gln

305		310		315		320
Lys Asn Leu Leu Phe Lys Asp Ser Thr Leu Glu Leu Val Pro Ile Ala						
		325		330		335
Thr Gln Thr Tyr Glu Ala Trp Leu Gly Pro Glu Tyr Leu His Ala Met						
		340		345		350
Lys Gly Leu Leu Cys Asp Pro Asn Arg Leu Pro Pro Tyr Leu Arg Trp						
		355		360		365
Cys Val Leu Ser Thr Pro Glu Ile Gln Lys Cys Gly Asp Met Ala Val						
		370		375		380
Ala Phe Ser Arg Gln Arg Leu Lys Pro Glu Ile Gln Cys Val Ser Ala						
		385		390		395
Glu Ser Pro Gln His Cys Met Glu Gln Ile Gln Ala Gly His Ile Asp						
		405		410		415
Ala Val Thr Leu Asn Gly Glu Asp Ile His Thr Ala Gly Lys Thr Tyr						
		420		425		430
Gly Leu Ile Pro Ala Ala Gly Glu Leu Tyr Ala Ala Asp Asp Arg Ser						
		435		440		445
Asn Ser Tyr Phe Val Val Ala Val Val Lys Arg Asp Ser Ala Tyr Ala						
		450		455		460
Phe Thr Val Asp Glu Leu Arg Gly Lys Arg Ser Cys His Pro Gly Phe						
		465		470		475
Gly Ser Pro Ala Gly Trp Asp Val Pro Val Gly Ala Leu Ile His Trp						
		485		490		495
Gly Tyr Ile Arg Pro Arg Asn Cys Asp Val Leu Thr Ala Val Gly Gln						
		500		505		510
Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Lys Tyr Pro						
		515		520		525
Ser Ser Leu Cys Ala Leu Cys Val Gly Asp Glu Gln Gly Arg Asn Lys						
		530		535		540
Cys Thr Gly Asn Ser Gln Glu Arg Tyr Tyr Gly Asp Ser Gly Ala Phe						
		545		550		555
Arg Cys Leu Val Glu Gly Ala Gly Asp Val Ala Phe Val Lys His Thr						
		565		570		575
Thr Ile Phe Asp Asn Thr Asn Gly His Asn Pro Glu Pro Trp Ala Ala						
		580		585		590
His Leu Arg Ser Gln Asp Tyr Glu Leu Leu Cys Pro Asn Gly Ala Arg						
		595		600		605
Ala Glu Ala His Gln Phe Ala Ala Cys Asn Leu Ala Gln Ile Pro Ser						
		610		615		620



His Ala Val Met Val Arg Pro Asp Thr Asn Ile Phe Thr Val Tyr Gly  
625 630 635 640

Leu Leu Asp Lys Ala Gln Asp Leu Phe Gly Asp Asp His Asn Lys Asn  
645 650 655

Gly Phe Lys Met Phe Asp Ser Ser Ser Tyr His Gly Arg Asp Leu Leu  
660 665 670

Phe Lys Asp Ala Thr Val Arg Ala Val Pro Val Gly Glu Arg Thr Thr  
675 680 685

Tyr Gln Asp Trp Leu Gly Pro Asp Tyr Val Ala Ala Leu Glu Gly Met  
690 695 700

Gln Ser Gln Arg Cys Ser Gly Ala Ala Val Gly Ala Pro Gly Ala Ser  
705 710 715 720

Leu Leu,Pro Leu Leu Pro Leu Ala Ala Gly Leu Leu Leu Ser Ser Leu  
725 730 735

```
<210> 3
<211> 2368
<212> DNA
<213> Homo sapiens
```

<400>	3						
gcgggacttcc	tcggaacccgg	accagacccc	agcccggccc	cagccagccc	cgacggcgcc		60
atgcgggggtc	cgagcggggc	tctgtggctg	ctcctggctc	tgcgcaccgt	gctcggaggc		120
atggaggtgc	ggtggtgctc	cacctcggac	ccagagcagc	acaagtgcgg	caacatgagc		180
gaggccttcc	gggaagcggg	catccagccc	tccctcctct	gcgtccgggg	cacctccgcc		240
gaccactgctg	tccagctcat	cgcggcccag	gaggctgacg	ccatcactct	ggatggagga		300
gccatctatg	aggcgggaaa	ggagcacggc	ctgaagccgg	tgggtgggcga	agtgtacgat		360
caagaggtcg	gtacctccta	ttacgcctg	gctgtggtca	ggaggagctc	ccatgtgacc		420
attgacaccc	tgaaaggcgt	gaagtctctg	cacacgggca	tcaatcgcac	agtgggctgg		480
aacgtgccc	tgggctacct	ggtggagagc	ggccgcctct	cggtgatggg	ctgcgatgta		540
ctcaaagctg	tcagcgacta	ttttgggggc	agctgcgtcc	cgggggcagg	agagaccagt		600
tactctgagt	ccctctgtcg	cctctgcagg	ggtgacagct	ctggggaagg	ggtgtgtgac		660
aagagcccc	tggagagata	ctacgactac	agcggggcct	tccggtgcct	ggcggaaggg		720
gcaggggacg	tggcttttgt	gaagcacagc	acggtactgg	agaacacgga	tgggaagacg		780
cttccctcct	ggggccaggc	cctgctgtca	caggacttcg	agctgctgtg	ccgggatggt		840
agccggggccg	atgtcaccca	gtggaggcag	tgccatctgg	cccgggtgcc	tgtcacgcc		900

gtggtggtcc gggccgacac agatgggggc ctcatcttcc ggctgctcaa cgaaggccag 960  
 cgtctgttca gccacgaggg cagcagcttc cagatgttca gctctgaggc ctatggccag 1020  
 aaggatctac tcttcaaaga ctctacctcg gagcttgtgc ccatcgccac acagacctat 1080  
 gaggcgtggc tgggccatga gtacctgcac gccatgaagg gtctgctctg tgaccccaac 1140  
 cggctgcccc cctacctgcg ctggtgtgtg ctctccactc ccgagatcca gaagtgtgga 1200  
 gacatggccg tggccttccg ccggcagcgc ctcaaggcag agatccagtg cgtgtcagcc 1260  
 aagtcccccc aacctgcat ggagcggatc caggctgagc aggtcgacgc tgtgacccta 1320  
 agtggcgagg acatttacac ggcggggaag aagtacggcc tggttcccg cagccggcgag 1380  
 cactatgccc cggaagacag cagcaactcg tactacgtgg tggccgtggt gagacgggac 1440  
 agctcccacg ccttcacctt ggatgagctt cggggcaagc gctcctgcca cgccggtttc 1500  
 ggcagccctg caggctggga tgtccccgtg ggtgccctta ttcagagagg cttcatccgg 1560  
 cccaaggact gtgacgtcct cacagcagtg agcgagttct tcaatgccag ctgctgccc 1620  
 gtgaacaacc ccaagaacta cccctcctcg ctgtgtgcac tgtgctggg ggacgagcag 1680  
 ggccgcaaca agtgtgtggg caacagccag gagcggattt acggctaccg cggcgccctc 1740  
 aggtgcctlg tggagaatgc gggtgacgtt gccttcgtca ggcacacaac cgtctttgac 1800  
 aacacaaacg gccacaattc cgagccctgg gctgctgagc tcaggtcaga ggactatgaa 1860  
 ctgctgtgcc ccaacggggc ccgagccgag gtgtcccagt ttgcagcctg caacctggca 1920  
 cagataccac cccacgccgt gatggtccgg cccgacacca acatcttcac cgtgtatgga 1980  
 ctgctggaca aggcccagga cctgttttga gacgaccaca ataagaacgg gttcaaaatg 2040  
 ttcgactcct ccaactatca tggccaagac ctgcttttca aggatgccac cgtccggggc 2100  
 gtgcctgtcg gagagaaaac cacctaccgc ggctggctgg ggctggacta cgtggcgggc 2160  
 ctggaaggga tgtcgtctca gcagtgtcg ggcgcagcgg ccccggcgc cggggcgccc 2220  
 ctgctcccg tgctgtgcc cgccctcgcc gccgcctgc tcccgccgc cctctgagcc 2280  
 cggccgcccc gcccagagc tccgatgcc gccgggggag ttccgcgggc ggccctctgc 2340  
 gctgcggaat ccagaaggaa gctcgca 2368

<210> 4  
 <211> 738  
 <212> PRT  
 <213> Homo sapiens  
 <400> 4

Met Arg Gly Pro Ser Gly Ala Leu Trp Leu Leu Leu Ala Leu Arg Thr  
 1 5 10 15  
 Val Leu Gly Gly Met Glu Val Arg Trp Cys Ala Thr Ser Asp Pro Glu  
 20 25 30  
 Gln His Lys Cys Gly Asn Met Ser Glu Ala Phe Arg Glu Ala Gly Ile  
 35 40 45  
 Gln Pro Ser Leu Leu Cys Val Arg Gly Thr Ser Ala Asp His Cys Val  
 50 55 60  
 Gln Leu Ile Ala Ala Gln Glu Ala Asp Ala Ile Thr Leu Asp Gly Gly  
 65 70 75 80  
 Ala Ile Tyr Glu Ala Gly Lys Glu His Gly Leu Lys Pro Val Val Gly  
 85 90 95  
 Glu Val Tyr Asp Gln Glu Val Gly Thr Ser Tyr Tyr Ala Val Ala Val  
 100 105 110  
 Val Arg Arg Ser Ser His Val Thr Ile Asp Thr Leu Lys Gly Val Lys  
 115 120 125  
 Ser Cys His Thr Gly Ile Asn Arg Thr Val Gly Trp Asn Val Pro Val  
 130 135 140  
 Gly Tyr Leu Val Glu Ser Gly Arg Leu Ser Val Met Gly Cys Asp Val  
 145 150 155 160  
 Leu Lys Ala Val Ser Asp Tyr Phe Gly Gly Ser Cys Val Pro Gly Ala  
 165 170 175  
 Gly Glu Thr Ser Tyr Ser Glu Ser Leu Cys Arg Leu Cys Arg Gly Asp  
 180 185 190  
 Ser Ser Gly Glu Gly Val Cys Asp Lys Ser Pro Leu Glu Arg Tyr Tyr  
 195 200 205  
 Asp Tyr Ser Gly Ala Phe Arg Cys Leu Ala Glu Gly Ala Gly Asp Val  
 210 215 220  
 Ala Phe Val Lys His Ser Thr Val Leu Glu Asn Thr Asp Gly Lys Thr  
 225 230 235 240  
 Leu Pro Ser Trp Gly Gln Ala Leu Leu Ser Gln Asp Phe Glu Leu Leu  
 245 250 255  
 Cys Arg Asp Gly Ser Arg Ala Asp Val Thr Glu Trp Arg Gln Cys His  
 260 265 270  
 Leu Ala Arg Val Pro Ala His Ala Val Val Val Arg Ala Asp Thr Asp  
 275 280 285  
 Gly Gly Leu Ile Phe Arg Leu Leu Asn Glu Gly Gln Arg Leu Phe Ser  
 290 295 300

His Glu Gly Ser Ser Phe Gln Met Phe Ser Ser Glu Ala Tyr Gly Gln  
305 310 315 320

Lys Asp Leu Leu Phe Lys Asp Ser Thr Ser Glu Leu Val Pro Ile Ala  
325 330 335

Thr Gln Thr Tyr Glu Ala Trp Leu Gly His Glu Tyr Leu His Ala Met  
340 345 350

Lys Gly Leu Leu Cys Asp Pro Asn Arg Leu Pro Pro Tyr Leu Arg Trp  
355 360 365

Cys Val Leu Ser Thr Pro Glu Ile Gln Lys Cys Gly Asp Met Ala Val  
370 375 380

Ala Phe Arg Arg Gln Arg Leu Lys Pro Glu Ile Gln Cys Val Ser Ala  
385 390 395 400

Lys Ser Pro Gln His Cys Met Glu Arg Ile Gln Ala Glu Gln Val Asp  
405 410 415

Ala Val Thr Leu Ser Gly Glu Asp Ile Tyr Thr Ala Gly Lys Lys Tyr  
420 425 430

Gly Leu Val Pro Ala Ala Gly Glu His Tyr Ala Pro Glu Asp Ser Ser  
435 440 445

Asn Ser Tyr Tyr Val Val Ala Val Val Arg Arg Asp Ser Ser His Ala  
450 455 460

Phe Thr Leu Asp Glu Leu Arg Gly Lys Arg Ser Cys His Ala Gly Phe  
465 470 475 480

Gly Ser Pro Ala Gly Trp Asp Val Pro Val Gly Ala Leu Ile Gln Arg  
485 490 495

Gly Phe Ile Arg Pro Lys Asp Cys Asp Val Leu Thr Ala Val Ser Glu  
500 505 510

Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Asn Tyr Pro  
515 520 525

Ser Ser Leu Cys Ala Leu Cys Val Gly Asp Glu Gln Gly Arg Asn Lys  
530 535 540

Cys Val Gly Asn Ser Gln Glu Arg Tyr Tyr Gly Tyr Arg Gly Ala Phe  
545 550 555 560

Arg Cys Leu Val Glu Asn Ala Gly Asp Val Ala Phe Val Arg His Thr  
565 570 575

Thr Val Phe Asp Asn Thr Asn Gly His Asn Ser Glu Pro Trp Ala Ala  
580 585 590

Glu Leu Arg Ser Glu Asp Tyr Glu Leu Leu Cys Pro Asn Gly Ala Arg  
595 600 605

Ala Glu Val Ser Gln Phe Ala Ala Cys Asn Leu Ala Gln Ile Pro Pro



```
<400> 7
agagggactc cgagtatctg gtctc
```

<210>	8
<211>	24
<212>	DNA
<213>	Artificial Sequence

<220>  
<223> synthetic

<400> 8  
gtccggcccg acaccaacat cttc 24

<210>	9
<211>	24
<212>	DNA
<213>	Artificial Sequence

<220>  
<223> synthetic

<400> 9  
ggtgtgttga ggggcgtgga ctct 24

<210>	10
<211>	25
<212>	DNA
<213>	Artificial Sequence

<220>  
<223> synthetic

```
<400> 10
tcaccaacgg ctttgagcac atcac                25
```

<210>	11
<211>	614
<212>	DNA
<213>	Mus sp.

[illegible]

gctccctgga aggcctcgct catgtctttg cacttctgct gctctgcgct tgagatggta 420  
 caccactgca cctccatcac acagacgaca gtgcgcaggg acaggagtag ccaaaaagtc 480  
 acgctcagga gcctcatggc aacgttgggt tggctggggg gctggcgggt ctgtcctggc 540  
 ttctcttccc ctggtctctc tggccttcac tattaagcg cagcccgggg agagtccacg 600  
 cccctcaaca cacc 614

<210> 12  
 <211> 17  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> synthetic

<400> 12 17  
 tgctacttca tcgaccc

<210> 13  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> synthetic

<400> 13 19  
 aaagacctcc cctccatct

<210> 14  
 <211> 4158  
 <212> DNA  
 <213> Mus sp.

<400> 14 60  
 ggtgtgttga ggggcgtgga ctctcccgg gctgcgctta aatagtgaag gccagagaga  
 ccagggaaga ggaagccagg acagaccgc cagcaccaca gccaaccaa cggtgccatg 120  
 aggctcctga gcgtgacttt ttggctactc ctgtccctgc gcactgtcgt ctgtgtgatg 180  
 gaggtgcagt ggtgtaccat ctcagacgca gagcagcaga agtgcaaaga catgagcgag 240  
 gccttccagg gagctggcat tcgtccttcc cttctctgcg tccagggcaa ctccgctgac 300  
 cactgtgtcc agctcatcaa ggaacaaaa gcagatgcca tcacctgga tggaggggcc 360  
 atctatgagg cagggaagga gcacggcctg aagccagtgg tgggggaagt ctatgaccaa 420  
 gacattggga cttcctatta tgccgtggct gtggtcagga ggaattccaa tgttaccatc 480







<210>	15
<211>	738
<212>	PRT
<213>	Mus sp.

<400> 15

Met 1	Arg	Leu	Leu	Ser 5	Val	Thr	Phe	Trp	Leu 10	Leu	Leu	Ser	Leu	Arg	Thr
Val	Val	Cys	Val	Met	Glu	Val	Gln	Trp 25	Cys	Thr	Ile	Ser	Asp	Ala	Glu
Gln	Gln	Lys 35	Cys	Lys	Asp	Met	Ser 40	Glu	Ala	Phe	Gln	Gly 45	Ala	Gly	Ile
Arg	Pro 50	Ser	Leu	Leu	Cys	Val 55	Gln	Gly	Asn	Ser	Ala 60	Asp	His	Cys	Val
Gln 65	Leu	Ile	Lys	Glu	Gln 70	Lys	Ala	Asp	Ala	Ile 75	Thr	Leu	Asp	Gly	Gly 80
Ala	Ile	Tyr	Glu	Ala 85	Gly	Lys	Glu	His	Gly 90	Leu	Lys	Pro	Val	Val	Gly
Glu	Val	Tyr	Asp 100	Gln	Asp	Ile	Gly	Thr 105	Ser	Tyr	Tyr	Ala	Val 110	Ala	Val
Val	Arg	Arg 115	Asn	Ser	Asn	Val	Thr 120	Ile	Asn	Thr	Leu	Lys 125	Gly	Val	Lys
Ser	Cys 130	His	Thr	Gly	Ile	Asn 135	Arg	Thr	Val	Gly	Trp 140	Asn	Val	Pro	Val
Gly 145	Tyr	Leu	Val	Glu	Ser 150	Gly	His	Leu	Ser	Val 155	Met	Gly	Cys	Asp	Val 160
Leu	Lys	Ala	Val	Gly 165	Asp	Tyr	Phe	Gly	Gly 170	Ser	Cys	Val	Pro	Gly 175	Thr
Gly	Glu	Thr	Ser 180	His	Ser	Glu	Ser	Leu 185	Cys	Arg	Leu	Cys	Arg 190	Gly	Asp
Ser	Ser	Gly 195	His	Asn	Val	Cys	Asp 200	Lys	Ser	Pro	Leu	Glu 205	Arg	Tyr	Tyr
Asp	Tyr 210	Ser	Gly	Ala	Phe	Arg 215	Cys	Leu	Ala	Glu	Gly 220	Ala	Gly	Asp	Val

Ala Phe Val Lys His Ser Thr Val Leu Glu Asn Thr Asp Gly Asn Thr  
225 230 235 240

Leu Pro Ser Trp Gly Lys Ser Leu Met Ser Glu Asp Phe Gln Leu Leu  
245 250 255

Cys Arg Asp Gly Ser Arg Ala Asp Ile Thr Glu Trp Arg Arg Cys His  
260 265 270

Leu Ala Lys Val Pro Ala His Ala Val Val Val Arg Gly Asp Met Asp  
275 280 285

Gly Gly Leu Ile Phe Gln Leu Leu Asn Glu Gly Gln Leu Leu Phe Ser  
290 295 300

His Glu Asp Ser Ser Phe Gln Met Phe Ser Ser Lys Ala Tyr Ser Gln  
305 310 315 320

Lys Asn Leu Leu Phe Lys Asp Ser Thr Leu Glu Leu Val Pro Ile Ala  
325 330 335

Thr Gln Asn Tyr Glu Ala Trp Leu Gly Gln Glu Tyr Leu Gln Ala Met  
340 345 350

Lys Gly Leu Leu Cys Asp Pro Asn Arg Leu Pro His Tyr Leu Arg Trp  
355 360 365

Cys Val Leu Ser Ala Pro Glu Ile Gln Lys Cys Gly Asp Met Ala Val  
370 375 380

Ala Phe Ser Arg Gln Asn Leu Lys Pro Glu Ile Gln Cys Val Ser Ala  
385 390 395 400

Glu Ser Pro Glu His Cys Met Glu Gln Ile Gln Ala Gly His Thr Asp  
405 410 415

Ala Val Thr Leu Arg Gly Glu Asp Ile Tyr Arg Ala Gly Lys Val Tyr  
420 425 430

Gly Leu Val Pro Ala Ala Gly Glu Leu Tyr Ala Glu Glu Asp Arg Ser  
435 440 445

Asn Ser Tyr Phe Val Val Ala Val Ala Arg Arg Asp Ser Ser Tyr Ser  
450 455 460

Phe Thr Leu Asp Glu Leu Arg Gly Lys Arg Ser Cys His Pro Tyr Leu  
465 470 475 480

Gly Ser Pro Ala Gly Trp Glu Val Pro Ile Gly Ser Leu Ile Gln Arg  
485 490 495

Gly Phe Ile Arg Pro Lys Asp Cys Asp Val Leu Thr Ala Val Ser Gln  
500 505 510

Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Asn Tyr Pro  
515 520 525

Ser Ala Leu Cys Ala Leu Cys Val Gly Asp Glu Lys Gly Arg Asn Lys  
530 535 540

Cys Val Gly Ser Ser Gln Glu Arg Tyr Tyr Gly Tyr Ser Gly Ala Phe  
545 550 555 560

Arg Cys Leu Val Glu His Ala Gly Asp Val Ala Phe Val Lys His Thr  
565 570 575

Thr Val Phe Glu Asn Thr Asn Gly His Asn Pro Glu Pro Trp Ala Ser  
580 585 590

His Leu Arg Trp Gln Asp Tyr Glu Leu Leu Cys Pro Asn Gly Ala Arg  
595 600 605

Ala Glu Val Asp Gln Phe Gln Ala Cys Asn Leu Ala Gln Met Pro Ser  
610 615 620

His Ala Val Met Val Arg Pro Asp Thr Asn Ile Phe Thr Val Tyr Gly  
625 630 635 640

Leu Leu Asp Lys Ala Gln Asp Leu Phe Gly Asp Asp His Asn Lys Asn  
645 650 655

Gly Phe Gln Met Phe Asp Ser Ser Lys Tyr His Ser Gln Asp Leu Leu  
660 665 670

Phe Lys Asp Ala Thr Val Arg Ala Val Pro Val Arg Glu Lys Thr Thr  
675 680 685

Tyr Leu Asp Trp Leu Gly Pro Asp Tyr Val Val Ala Leu Glu Gly Met  
690 695 700

Leu Ser Gln Gln Cys Ser Gly Ala Gly Ala Ala Val Gln Arg Val Pro  
705 710 715 720

Leu Leu Ala Leu Leu Leu Leu Thr Leu Ala Ala Gly Leu Leu Pro Arg  
725 730 735

Val Leu

Sequence Listing

<110> CHUGAI SEIYAKU KABUSHIKI KAISHA  
 <120> CHONDROGENESIS STIMULATOR  
 5 <130> YCT-501  
 <160> 15  
  
 <210> 1  
 <211> 2388  
 10 <212> cDNA  
 <213> Rabbit  
 <400> 1  
 gccgccgctc actcggtcgc actcggactc agaccagtc cgacccctg gactgcgcca 60  
 tgcgggtgcg aagcgcggct atgtggatct tcctggccct gcgcaccgca ctcggcagcg 120  
 15 tggaggtgcg gtggtgcacc gcgtccgagc ccgagcagca gaagtgcgag gacatgagcc 180  
 aggccttccg cgaagccggc ctccagcccg cctgtctgtg cgtgcagggc acctcggccg 240  
 accactgcgt ccagctcatc gcggcccacg aggcgcagc catcactctg gacggaggag 300  
 ccatttaaga ggcggggaag gaacacggcc tgaagcccggt ggtgggcgaa gtgtatgacc 360  
 aagaggtggg cacctcctac tacgtgtgg ccgtggtcaa gaggagctcc aacgtgacca 420  
 20 tcaacaccct gagaggcgtg aagtcctgcc acacgggcat caaccgcacg gtgggctgga 480  
 acgtgcctgt gggctacctg gtggacagcg gccgcctctc agtgatgggc tgtgacgtgc 540  
 tcaaagcggc cagcgagtac ttcgggggca gctgcgtccc tggggcagga gagaccagat 600  
 actcggagtc cctctgtcgc ctctgcggg gcgacacctc cggggagggg gtgtgtgaca 660  
 agagccccct ggagcggtag tacgactaca gcggggcctt ccggtgcctg gcagaaggcg 720  
 25 caggggacgt ggcctttgtg aagcacagca cgggtgctga gaacacggat gggagaacac 780  
 tgccctcctg gggccacatg ctgatgtcac gggactttga gctgctgtgc cgggacggca 840  
 gccgggccag cgtcaccgag tggcagcact gccacctggc ccgggtgccc gccacggcg 900  
 tgggtggtccg ggccgacacc gacgcaggcc tcattctccg gcttctcaat gagggccagc 960  
 ggctgttcag ccacgagggc agcagcttcc agatgttcag ctcgagggcc tacggccaga 1020

agaacctgct gttcaaagac tccacgctgg agctggtgcc catcgccaca cagacctacg 1080  
 aggcctggct gggccccgag tacctgcacg ccatgaaggg tctgctctgt gacccaacc 1140  
 ggctgcccc atacctgccc tgggtgcgtgc tgtccacccc cgagatccag aagtgtggag 1200  
 acatggccgt ggccttcagc cggcagaggc tcaagccgga gatccagtgt gtctcggcgg 1260  
 5 agtcccccca gcactgcatg gagcagatcc aggctgggca catcgatgct gtgaccctga 1320  
 acggggagga cattcacaca gcggggaaga cttatgggct gatcccggt gccggggagc 1380  
 tgtatgccgc ggacgacagg agtaactcgt acttcgtggt ggccgtggtg aagcgagaca 1440  
 ggcctacgc cttaccgtg gacgagctgc gcgggaagcg ctctgccac cccggttcg 1500  
 gcagcccggc cggctgggac gtcccgttg gcgccctcat ccaactggggc tacatccggc 1560  
 10 ccaggaactg cgacgtcctc acagcggtag gtcagttctt caacgccagc tgtgtgccgg 1620  
 tgaacaacc caagaagtac ccctcctcgc tgtgcgcact ctgcgtgggt gacgagcagg 1680  
 gccgcaacaa gtgcactggc aacagccagg agcggtaacta tggcgacagt ggcgccttca 1740  
 ggtgcctggt ggagggtgca ggggacgtgg ccttcgtcaa gcacacgacc atctttgaca 1800  
 acacaaatgg ccacaatccc gagccgtggg ctgccatct gaggagccag gactacgagc 1860  
 15 tgctgtgccc caacggcgcg cgagctgagg cgcaccagtt tgccgcctgc aacctggccc 1920  
 agattccgtc ccacgccgtc atggtgcggc ccgacaccaa catcttcacc gtttacggac 1980  
 tgctggacaa ggcccaggac ctgtttggag acgaccacaa caagaacggg ttcaagatgt 2040  
 tcgactcctc cagctaccac ggccgagacc tgctcttcaa ggacgccag gtgcgcgctg 2100  
 tgctgtggg cgagaggacc acctaccagg actggctggg gccggactac gtggcggctc 2160  
 20 tggaagggat gcagtcacag cgggtgctcag gggcagccgt cggcgcccc ggggcctcgc 2220  
 tgctgccgt gctgccctg gctgcgggccc tctgctgtc ttcgctctga gacgagcccc 2280  
 gggcagcctc gggcccgga ggggagcctg cgcggaagct tctgaacga gcccgcgccc 2340  
 tggctggatg tggttacctc ggcgagccgc ggggcgcgc tcccccg 2388

- 25 <210> 2  
 <211> 736  
 <212> PRT  
 <213> Rabbit  
 <400> 2

$$\begin{aligned} & \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \\ & \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) \end{aligned}$$

	Met	Arg	Cys	Arg	Ser	Ala	Ala	Met	Trp	Ile	Phe	Leu	Ala	Leu	Arg	Thr
	1				5					10					15	
	Ala	Leu	Gly	Ser	Val	Glu	Val	Arg	Trp	Cys	Thr	Ala	Ser	Glu	Pro	Glu
				20					25					30		
5	Gln	Gln	Lys	Cys	Glu	Asp	Met	Ser	Gln	Ala	Phe	Arg	Glu	Ala	Gly	Leu
			35					40					45			
	Gln	Pro	Ala	Leu	Leu	Cys	Val	Gln	Gly	Thr	Ser	Ala	Asp	His	Cys	Val
		50					55					60				
	Gln	Leu	Ile	Ala	Ala	His	Glu	Ala	Asp	Ala	Ile	Thr	Leu	Asp	Gly	Gly
10	65					70					75				80	
	Ala	Ile	Tyr	Glu	Ala	Gly	Lys	Glu	His	Gly	Leu	Lys	Pro	Val	Val	Gly
					85					90				95		
	Glu	Val	Tyr	Asp	Gln	Glu	Val	Gly	Thr	Ser	Tyr	Tyr	Ala	Val	Ala	Val
				100					105					110		
15	Val	Lys	Arg	Ser	Ser	Asn	Val	Thr	Ile	Asn	Thr	Leu	Arg	Gly	Val	Lys
			115					120					125			
	Ser	Cys	His	Thr	Gly	Ile	Asn	Arg	Thr	Val	Gly	Trp	Asn	Val	Pro	Val
		130					135					140				
	Gly	Tyr	Leu	Val	Asp	Ser	Gly	Arg	Leu	Ser	Val	Met	Gly	Cys	Asp	Val
20	145					150					155				160	
	Leu	Lys	Ala	Val	Ser	Glu	Tyr	Phe	Gly	Gly	Ser	Cys	Val	Pro	Gly	Ala
				165					170					175		
	Gly	Glu	Thr	Arg	Tyr	Ser	Glu	Ser	Leu	Cys	Arg	Leu	Cys	Arg	Gly	Asp
			180						185					190		
25	Thr	Ser	Gly	Glu	Gly	Val	Cys	Asp	Lys	Ser	Pro	Leu	Glu	Arg	Tyr	Tyr
			195						200					205		
	Asp	Tyr	Ser	Gly	Ala	Phe	Arg	Cys	Leu	Ala	Glu	Gly	Ala	Gly	Asp	Val
		210					215						220			
	Ala	Phe	Val	Lys	His	Ser	Thr	Val	Leu	Glu	Asn	Thr	Asp	Gly	Arg	Thr

	225		230		235		240									
	Leu	Pro	Ser	Trp	Gly	His	Met	Leu	Met	Ser	Arg	Asp	Phe	Glu	Leu	Leu
					245					250					255	
	Cys	Arg	Asp	Gly	Ser	Arg	Ala	Ser	Val	Thr	Glu	Trp	Gln	His	Cys	His
5				260						265					270	
	Leu	Ala	Arg	Val	Pro	Ala	His	Ala	Val	Val	Val	Arg	Ala	Asp	Thr	Asp
				275						280					285	
	Ala	Gly	Leu	Ile	Phe	Arg	Leu	Leu	Asn	Glu	Gly	Gln	Arg	Leu	Phe	Ser
				290						295					300	
10	His	Glu	Gly	Ser	Ser	Phe	Gln	Met	Phe	Ser	Ser	Glu	Ala	Tyr	Gly	Gln
				305						310					315	
	Lys	Asn	Leu	Leu	Phe	Lys	Asp	Ser	Thr	Leu	Glu	Leu	Val	Pro	Ile	Ala
										325					330	
	Thr	Gln	Thr	Tyr	Glu	Ala	Trp	Leu	Gly	Pro	Glu	Tyr	Leu	His	Ala	Met
15				340						345					350	
	Lys	Gly	Leu	Leu	Cys	Asp	Pro	Asn	Arg	Leu	Pro	Pro	Tyr	Leu	Arg	Trp
				355						360					365	
	Cys	Val	Leu	Ser	Thr	Pro	Glu	Ile	Gln	Lys	Cys	Gly	Asp	Met	Ala	Val
				370						375					380	
20	Ala	Phe	Ser	Arg	Gln	Arg	Leu	Lys	Pro	Glu	Ile	Gln	Cys	Val	Ser	Ala
				385						390					395	
	Glu	Ser	Pro	Gln	His	Cys	Met	Glu	Gln	Ile	Gln	Ala	Gly	His	Ile	Asp
										405					410	
	Ala	Val	Thr	Leu	Asn	Gly	Glu	Asp	Ile	His	Thr	Ala	Gly	Lys	Thr	Tyr
25				420						425					430	
	Gly	Leu	Ile	Pro	Ala	Ala	Gly	Glu	Leu	Tyr	Ala	Ala	Asp	Asp	Arg	Ser
				435						440					445	
	Asn	Ser	Tyr	Phe	Val	Val	Ala	Val	Val	Lys	Arg	Asp	Ser	Ala	Tyr	Ala
				450						455					460	







gaggcgtggc tgggcatga gtacctgcac gccatgaagg gtctgctctg tgaccccaac 1140  
 cggctgcccc cctacctgcg ctggtgtgtg ctctccactc ccgagatcca gaagtgtgga 1200  
 gacatggccg tggccttcg ccggcagcgc ctcaagccag agatccagtg cgtgtcagcc 1260  
 aagtcccccc aacctgcat ggagcggatc caggctgagc aggtcgacgc tgtgacccta 1320  
 5 agtggcgagg acatttacac ggcggggaag aagtacggcc tggttccgc agccggcgag 1380  
 cactatgccc cggaagacag cagcaactcg tactacgtgg tggcgtggt gagacgggac 1440  
 agtccccacg cttcacctt ggatgagctt cggggcaagc gctcctgcca cgccggtttc 1500  
 ggcagccctg caggctggga tgtccccgtg ggtgccctta ttcagagagg cttcatccgg 1560  
 cccaaggact gtgacgtcct cacagcagtg agcgagttct tcaatgccag ctgctgccc 1620  
 10 gtgaacaacc ccaagaacta cccctcctcg ctgtgtgcac tgtgctggg ggacgagcag 1680  
 ggccgcaaca agtgtgtggg caacagccag gagcggattt acggctaccg cggcgccttc 1740  
 aggtgcctgg tggagaatgc gggtgacgtt gccttcgtca ggcacacaac cgtctttgac 1800  
 aacacaaacg gccacaattc cgagccctgg gctgctgagc tcaggtcaga ggactatgaa 1860  
 ctgctgtgcc ccaacggggc ccgagccgag gtgtcccagt ttgcagcctg caacctggca 1920  
 15 cagataccac cccacgccgt gatggtccg cccgacacca acatcttcac cgtgtatgga 1980  
 ctgctggaca aggcccagga cctgtttgga gacgaccaca ataagaacgg gttcaaaatg 2040  
 ttcgactcct ccaactatca tggccaagac ctgcttttca aggatgccac cgtccggggc 2100  
 gtgcctgtcg gagagaaaac cacctaccgc ggctggctgg ggctggacta cgtggcggcg 2160  
 ctggaaggga tgtcgtctca gcagtgtcg ggcgcagcgg ccccggcgcc cggggcgccc 2220  
 20 ctgctccgc tgctgtgcc cgccctgcgc gccgcctgc tcccgccgc cctctgagcc 2280  
 cggccgcccc gccccagagc tccgatgcc gcccggggag tttccgcggc ggcctctgc 2340  
 gctgcggaat ccagaaggaa gctcgca 2368

<210> 4

25 <211> 738

<212> PRT

<213> Human

<400> 4

Met Arg Gly Pro Ser Gly Ala Leu Trp Leu Leu Leu Ala Leu Arg Thr

	1	5	10	15
	Val	Leu	Gly	Gly
	Met	Glu	Val	Arg
	Trp	Cys	Ala	Thr
	Ser	Asp	Pro	Glu
	20	25	30	
	Gln	His	Lys	Cys
	Gly	Asn	Met	Ser
	Glu	Ala	Phe	Arg
	Glu	Ala	Gly	Ile
5	35	40	45	
	Gln	Pro	Ser	Leu
	Leu	Cys	Val	Arg
	Gly	Thr	Ser	Ala
	Asp	His	Cys	Val
	50	55	60	
	Gln	Leu	Ile	Ala
	Ala	Gln	Glu	Ala
	Asp	Ala	Ile	Thr
	Leu	Asp	Gly	Gly
	65	70	75	80
10	Ala	Ile	Tyr	Glu
	Ala	Gly	Lys	Glu
	His	Gly	Leu	Lys
	Pro	Val	Val	Gly
	85	90	95	
	Glu	Val	Tyr	Asp
	Gln	Glu	Val	Gly
	Thr	Ser	Tyr	Tyr
	Ala	Val	Ala	Val
	100	105	110	
	Val	Arg	Arg	Ser
	Ser	His	Val	Thr
	Ile	Asp	Thr	Leu
	Lys	Gly	Val	Lys
15	115	120	125	
	Ser	Cys	His	Thr
	Gly	Ile	Asn	Arg
	Thr	Val	Gly	Trp
	Asn	Val	Pro	Val
	130	135	140	
	Gly	Tyr	Leu	Val
	Glu	Ser	Gly	Arg
	Leu	Ser	Val	Met
	Gly	Cys	Asp	Val
	145	150	155	160
20	Leu	Lys	Ala	Val
	Ser	Asp	Tyr	Phe
	Gly	Gly	Ser	Cys
	Val	Pro	Gly	Ala
	165	170	175	
	Gly	Glu	Thr	Ser
	Tyr	Ser	Glu	Ser
	Leu	Cys	Arg	Leu
	Cys	Arg	Gly	Asp
	180	185	190	
	Ser	Ser	Gly	Glu
	Gly	Val	Cys	Asp
	Lys	Ser	Pro	Leu
	Glu	Arg	Tyr	Tyr
25	195	200	205	
	Asp	Tyr	Ser	Gly
	Ala	Phe	Arg	Cys
	Leu	Ala	Glu	Gly
	Ala	Gly	Asp	Val
	210	215	220	
	Ala	Phe	Val	Lys
	His	Ser	Thr	Val
	Leu	Glu	Asn	Thr
	Asp	Gly	Lys	Thr
	225	230	235	240



	465	470	475	480
	Gly Ser Pro Ala Gly Trp Asp Val Pro Val Gly Ala Leu Ile Gln Arg			
	485	490	495	
	Gly Phe Ile Arg Pro Lys Asp Cys Asp Val Leu Thr Ala Val Ser Glu			
5	500	505	510	
	Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Asn Tyr Pro			
	515	520	525	
	Ser Ser Leu Cys Ala Leu Cys Val Gly Asp Glu Gln Gly Arg Asn Lys			
	530	535	540	
10	Cys Val Gly Asn Ser Gln Glu Arg Tyr Tyr Gly Tyr Arg Gly Ala Phe			
	545	550	555	560
	Arg Cys Leu Val Glu Asn Ala Gly Asp Val Ala Phe Val Arg His Thr			
	565	570	575	
	Thr Val Phe Asp Asn Thr Asn Gly His Asn Ser Glu Pro Trp Ala Ala			
15	580	585	590	
	Glu Leu Arg Ser Glu Asp Tyr Glu Leu Leu Cys Pro Asn Gly Ala Arg			
	595	600	605	
	Ala Glu Val Ser Gln Phe Ala Ala Cys Asn Leu Ala Gln Ile Pro Pro			
	610	615	620	
20	His Ala Val Met Val Arg Pro Asp Thr Asn Ile Phe Thr Val Tyr Gly			
	625	630	635	640
	Leu Leu Asp Lys Ala Gln Asp Leu Phe Gly Asp Asp His Asn Lys Asn			
	645	650	655	
	Gly Phe Lys Met Phe Asp Ser Ser Asn Tyr His Gly Gln Asp Leu Leu			
25	660	665	670	
	Phe Lys Asp Ala Thr Val Arg Ala Val Pro Val Gly Glu Lys Thr Thr			
	675	680	685	
	Tyr Arg Gly Trp Leu Gly Leu Asp Tyr Val Ala Ala Leu Glu Gly Met			
	690	695	700	

Ser Ser Gln Gln Cys Ser Gly Ala Ala Ala Pro Ala Pro Gly Ala Pro  
 705 710 715 720  
 Leu Leu Pro Leu Leu Leu Pro Ala Leu Ala Ala Arg Leu Leu Pro Pro  
 725 730 735

5 Ala Leu

<210> 5

<211> 23

<212> DNA

10 <213> Artificial Sequence

<400> 5

ggctggaacg tgcccgtggg cta

<210> 6

15 <211> 23

<212> DNA

<213> Artificial Sequence

<400> 6

gtcctggggcc ttgtccagca gtc

20

<210> 7

<211> 25

<212> DNA

<213> Artificial Sequence

25 <400> 7

agagggactc cgagtatctg gtctc

<210> 8

<211> 24

<212> DNA  
 <213> Artificial Sequence  
 <400> 8  
 gtccggcccc acaccaacat ctct  
 5  
 <210> 9  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence  
 10 <400> 9  
 ggtgtgttga ggggcgtgga ctct  
  
 <210> 10  
 <211> 25  
 15 <212> DNA  
 <213> Artificial Sequence  
 <400> 10  
 tcaccaacgg ctttgagcac atcac  
  
 20 <210> 11  
 <211> 614  
 <212> RNA  
 <213> Mouse  
 <400> 11  
 25 tcaccaacgg ctttgagcac atcacagccc atcactgaca gatggccgct ctctacgagg 60  
 taaccgacag gcacgttcca gccacagtc cggttaatgc ctgtgtggca ggacttgacg 120  
 cccttcaggg tgttgatggt aacattggaa ttctcctga ccacagccac ggcataatag 180  
 gaagtcccaa tgtcttggtc atagacttcc cccaccactg gcttcaggcc gtgctccttc 240  
 cctgcctcat agatggcccc tccatccagg gtgatggcat ctgctttttg ttcttgatg 300



agctggacac agtggtcagc ggagttgccc tggacgcaga gaaggggaagg acgaatgcca 360  
gctccctgga aggcctcgct catgtctttg cacttctgct gctctgcgct tgagatggta 420  
caccactgca cctccatcac acagacgaca gtgcgcaggg acaggagtag ccaaaaagtc 480  
acgctcagga gcctcatggc aacgttgggt tggctggggg gctggcgggg ctgtcctggc 540  
5 ttctcttcc ctggtctctc tggccttcac tatttaagcg cagcccgggg agagtccacg 600  
ccctcaaca cacc 614

<210> 12  
<211> 17  
10 <212> DNA  
<213> Artificial Sequence  
<400> 12  
tgctacttca tcgaccc

15 <210> 13  
<211> 19  
<212> DNA  
<213> Artificial Sequence  
<400> 13

20 aaagacctcc cctccatct

<210> 14  
<211> 4158  
<212> cDNA  
25 <213> Mouse  
<400> 14

ggtgtgttga ggggcgtgga ctctccccgg gctgcgctta aatagtgaag gccagagaga 60  
ccaggggaaga ggaagccagg acagacccgc cagcacccca gccaaaccaa cgttgccatg 120  
aggctcctga gcgtgacttt ttggctactc ctgtccctgc gcaactgtcgt ctgtgtgatg 180

	gaggtgcagt ggtgtacat ctcagacgca gagcagcaga agtgcaaaga catgagcgag	240
	gccttcagg gagctggcat tcgtccttcc ctctctctgcg tccagggcaa ctccgctgac	300
	caactgtgtcc agctcatcaa ggaacaaaaa gcagatgccca tcacctgga tggagggggcc	360
	atctatgagg caggggaagga gcacggcctg aagccagtgg tgggggaagt ctatgaccaa	420
5	gacattggga ctctctatta tgccgtggct gtggtcagga ggaattccaa tgttaccatc	480
	aacaccctga agggcgtaaa gtcttgccac acaggcatta accggactgt gggctggaac	540
	gtgcctgtcg gttacctcgt agagagcggc catctgtcag tgatgggctg tgatgtgctc	600
	aaagccgttg gtgattatct tggaggcagc tgtgtccctg gaacaggaga aaccagccat	660
	tccgagtcct tctgtgcct ctgccgtggc gactcttctg ggcacaatgt gtgtgacaag	720
10	agtcccctag agagatacta cgactacagt ggagccttcc ggtgcctggc ggaaggagcc	780
	ggtgacgtgg ccttcgtgaa gcacagcaca gtgctggaaa atactgatgg aaacaccctg	840
	ccttcctggg gcaagtcctt gatgtcagag gaattccagc tactatgcag ggatggcagc	900
	cgagccgaca tcaactgagt gagacgttgc cacctggcca aggtgcctgc tcatgctgtg	960
	gtggtcaggg gtgacatgga tggcgggtctc atattccaac tgctcaacga aggccagctt	1020
15	ctgttcagcc acgaagacag cagcttccag atgttcagct ccaaagccta cagccagaag	1080
	aaacttgctgt tcaaagactc caccttggag cttgtgcccc ttgccacaca gaactacgag	1140
	gcctggctgg gccaggaata cctgcaggcc atgaaggggc tcctctgtga tccaaccgg	1200
	ctgccccact acctgcgctg gtgtgtgctg tcagcgcccg agatccagaa gtgtggagat	1260
	atggctgtgg ccttcagccg ccagaatctc aagccggaaa ttcagtgtgt gtcggccgag	1320
20	tcccctgagc actgcatgga gcagatccag gctgggcaca ctgacgctgt gactctgagg	1380
	ggcgaggaca ttacagggc aggaaagggt tacggcctgg ttccggcggc cggggagctg	1440
	tatgctgagg aggacaggag caattcctac tttgtgggtg ctgtggcaag aagggacagc	1500
	tcctactcct tcacctgga cgagcttcgc ggcaagcggt cctgccaccc ctacttgggc	1560
	agcccagcgg gctgggaggt gcccatcggc tccctcatcc agcggggctt catccggccc	1620
25	aaggactgtg atgtcctcac agcggtgagc cagttcttca atgccagctg cgtgcctgtc	1680
	aacaacccta agaactaccc ttccgcacta tgtgcgctct gcgtgggaga cgagaagggc	1740
	cgcaacaaat gtgtggggag cagccaggag agatactacg gctacagcgg ggccttcagg	1800
	tgcccttggtg agcatgcagg ggacgtggct ttcgtcaagc acacgactgt ctttgagaac	1860
	acaaatgggc acaatcctga gccttgggtt tctcacctca ggtggcaaga ctatgaacta	1920



ggccgggagc tatttgtgtc ttggtctgtg tggggggcct cagatcccaa tgacaggcca 3720  
 ggttcccagt ggctcgcccc cacctgtggg cgacgacggg acagatcctt tccatggctc 3780  
 accagtagag aaggtcctgg cagtgtccca gccagagtca cacaatcctg aggaaaatcg 3840  
 gtcaccatgg tgcttgggag agcaagcccc tcctcctccc agtacacagc catccattct 3900  
 5 tctctgagct ggggacttca cagtgagaag tgtactctgt gtgggcgact gtgctgcca 3960  
 aagtgtgatg tctgtgccgt gtgcctttca ggtgtgactt tgaagagcgt tgtgtaaagt 4020  
 acgtctgatt gccatgggcc actgctgtgt ttgtgctaaa gaaagacatt ggtttctttt 4080  
 taaaataaag ccatatatcc ctgcatacgc agaggcttgg atcctgggtg aaaaaaaaaa 4140  
 aaaaaaaaaa aaaaaaaaaa 4158

10

<210> 15  
 <211> 737  
 <212> PRT  
 <213> Mouse

15

<400> 15  
 Met Arg Leu Leu Ser Val Thr Phe Trp Leu Leu Leu Ser Leu Arg Thr  
 1 5 10 15  
 Val Val Cys Val Met Glu Val Gln Trp Cys Thr Ile Ser Asp Ala Glu  
 20 25 30  
 20 Gln Gln Lys Cys Lys Asp Met Ser Glu Ala Phe Gln Gly Ala Gly Ile  
 35 40 45  
 Arg Pro Ser Leu Leu Cys Val Gln Gly Asn Ser Ala Asp His Cys Val  
 50 55 60  
 Gln Leu Ile Lys Glu Gln Lys Ala Asp Ala Ile Thr Leu Asp Gly Gly  
 25 65 70 75 80  
 Ala Ile Tyr Glu Ala Gly Lys Glu His Gly Leu Lys Pro Val Val Gly  
 85 90 95  
 Glu Val Tyr Asp Gln Asp Ile Gly Thr Ser Tyr Tyr Ala Val Ala Val  
 100 105 110



	340	345	350
	Lys Gly Leu Leu Cys Asp Pro Asn Arg Leu Pro His Tyr Leu Arg Trp		
	355	360	365
	Cys Val Leu Ser Ala Pro Glu Ile Gln Lys Cys Gly Asp Met Ala Val		
5	370	375	380
	Ala Phe Ser Arg Gln Asn Leu Lys Pro Glu Ile Gln Cys Val Ser Ala		
	385	390	395 400
	Glu Ser Pro Glu His Cys Met Glu Gln Ile Gln Ala Gly His Thr Asp		
	405	410	415
10	Ala Val Thr Leu Arg Gly Glu Asp Ile Tyr Arg Ala Gly Lys Val Tyr		
	420	425	430
	Gly Leu Val Pro Ala Ala Gly Glu Leu Tyr Ala Glu Glu Asp Arg Ser		
	435	440	445
	Asn Ser Tyr Phe Val Val Ala Val Ala Arg Arg Asp Ser Ser Tyr Ser		
15	450	455	460
	Phe Thr Leu Asp Glu Leu Arg Gly Lys Arg Ser Cys His Pro Tyr Leu		
	465	470	475 480
	Gly Ser Pro Ala Gly Trp Glu Val Pro Ile Gly Ser Leu Ile Gln Arg		
	485	490	495
20	Gly Phe Ile Arg Pro Lys Asp Cys Asp Val Leu Thr Ala Val Ser Gln		
	500	505	510
	Phe Phe Asn Ala Ser Cys Val Pro Val Asn Asn Pro Lys Asn Tyr Pro		
	515	520	525
	Ser Ala Leu Cys Ala Leu Cys Val Gly Asp Glu Lys Gly Arg Asn Lys		
25	530	535	540
	Cys Val Gly Ser Ser Gln Glu Arg Tyr Tyr Gly Tyr Ser Gly Ala Phe		
	545	550	555 560
	Arg Cys Leu Val Glu His Ala Gly Asp Val Ala Phe Val Lys His Thr		
	565	570	575



Thr Val Phe Glu Asn Thr Asn Gly His Asn Pro Glu Pro Trp Ala Ser  
580 585 590  
His Leu Arg Trp Gln Asp Tyr Glu Leu Leu Cys Pro Asn Gly Ala Arg  
595 600 605  
5 Ala Glu Val Asp Gln Phe Gln Ala Cys Asn Leu Ala Gln Met Pro Ser  
610 615 620  
His Ala Val Met Val Arg Pro Asp Thr Asn Ile Phe Thr Val Tyr Gly  
625 630 635 640  
Leu Leu Asp Lys Ala Gln Asp Leu Phe Gly Asp Asp His Asn Lys Asn  
10 645 650 655  
Gly Phe Gln Met Phe Asp Ser Ser Lys Tyr His Ser Gln Asp Leu Leu  
660 665 670  
Phe Lys Asp Ala Thr Val Arg Ala Val Pro Val Arg Glu Lys Thr Thr  
675 680 685  
15 Tyr Leu Asp Trp Leu Gly Pro Asp Tyr Val Val Ala Leu Glu Gly Met  
690 695 700  
Leu Ser Gln Gln Cys Ser Gly Ala Gly Ala Ala Val Gln Arg Val Pro  
705 710 715 720  
Leu Leu Ala Leu Leu Leu Leu Thr Leu Ala Ala Gly Leu Leu Pro Arg  
20 725 730 735  
Val Leu

## Combined Declaration for Patent Application and Power of Attorney

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

**CHONDROGENESIS STIMULATOR**

the specification of which (check one)

- ☐ is attached hereto;  
☐ was filed in the United States under 35 U.S.C. §111 on \_\_\_\_\_, as  
 U.S. Appln. No. \_\_\_\_\_\*; or  
☒ was/will be filed in the U.S. under 35 U.S.C. §371 by entry into the U.S. national stage of an international (PCT) application, PCT/JP00/05590 filed Aug. 21, 2000, entry requested on \_\_\_\_\_\*; national stage application received U.S. Appln. No. \_\_\_\_\_\*, §371/§102(e) date \_\_\_\_\_\* (\* if known)

and was amended on \_\_\_\_\_ (if applicable).

*(include dates of amendments under PCT Art. 19 and 34 if PCT)*

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and I acknowledge the duty to disclose to the Patent and Trademark Office (PTO) all information known by me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119 and 365 of any prior foreign application(s) for patent or inventor's certificate, or prior PCT application(s) designating a country other than the U.S., listed below with the "Yes" box checked and have also identified below any such application having a filing date before that of the application on which priority is claimed:

<u>232966/1999</u>	<u>Japan</u>	<u>19/8/1999</u>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day Month Year Filed)	YES	NO
_____	_____	_____	<input type="checkbox"/>	<input type="checkbox"/>
(Number)	(Country)	(Day Month Year Filed)	YES	NO

I hereby claim the benefit under 35 U.S.C. §120 of any prior U.S. non-provisional application(s) or prior PCT application(s) designating the U.S. listed below, or under §119(e) of any prior U.S. provisional applications listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in such U.S. or PCT application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the PTO all information as defined in 37 C.F.R. §1.56(a) which occurred between the filing date of the prior application and the national filing date of this application:

_____ (Application No.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)
_____ (Application No.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)
_____ (Application No.)	_____ (Day Month Year Filed)	_____ (Status: patented, pending, abandoned)

As a named inventor, I hereby appoint the following registered practitioners to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

**All of the practitioners associated with Customer Number 001444**

Direct all correspondence to the address associated with Customer Number 001444; i.e.,  
**BROWDY AND NEIMARK, P.L.L.C.**  
 624 Ninth Street, N.W.  
 Washington, D.C. 20001-5303  
 (202) 628-5197

The undersigned hereby authorizes the U.S. Attorneys or Agents appointed herein to accept and follow instructions from YUASA AND HARA as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorneys or Agents and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents appointed herein will be so notified by the undersigned.



Page 2 of 2 Pages

Atty. Docket:

Title: CHONDROGENESIS STIMULATOR

U.S. Application filed \_\_\_\_\_, Serial No. \_\_\_\_\_

PCT Application filed August 21, 2000, Serial No. PCT/JP00/05590

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF FIRST INVENTOR <u>Yukio KATO</u>		INVENTOR'S SIGNATURE <u>Yukio Kato</u>	DATE <u>Feb. 1, 2002</u>
RESIDENT <u>Hiroshima, Japan</u> <i>gp x</i>		CITIZENSHIP <u>Japanese</u>	
POST OFFICE ADDRESS <u>3-6-9-501, Ushitawaseda, Higashi-ku, Hiroshima-shi,</u> <u>Hiroshima 732-0062 Japan</u>			
FULL NAME OF SECOND JOINT INVENTOR <u>Katsumi FUJIMOTO</u>		INVENTOR'S SIGNATURE <u>Katsumi Fujimoto</u>	DATE <u>Feb. 1, 2002</u>
RESIDENT <u>Hiroshima, Japan</u> <i>gp x</i>		CITIZENSHIP <u>Japanese</u>	
POST OFFICE ADDRESS <u>1-3-11-202, Asahi, Minami-ku, Hiroshima-shi,</u> <u>Hiroshima 734-0036 Japan</u>			
FULL NAME OF THIRD JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF FOURTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF FIFTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF SIXTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			
FULL NAME OF SEVENTH JOINT INVENTOR		INVENTOR'S SIGNATURE	DATE
RESIDENT		CITIZENSHIP	
POST OFFICE ADDRESS			

ALL INVENTORS MUST REVIEW APPLICATION AND DECLARATION BEFORE SIGNING. ALL ALTERATIONS MUST BE INITIALED AND DATED BY ALL INVENTORS PRIOR TO EXECUTION. NO ALTERATIONS CAN BE MADE AFTER THE DECLARATION IS SIGNED. ALL PAGES OF DECLARATION MUST BE SEEN BY ALL INVENTORS.

INVENTOR INFORMATION

10/049957  
JC13 Rec'd PCT/PTO 19 FEB 2002

Inventor One Given Name:: Yukio  
Family Name:: KATO  
Postal Address Line One:: 3-6-9-501, Ushitawaseda, Higashi-ku  
Postal Address Line Two:: Hiroshima-shi  
City:: Hiroshima  
Country:: Japan  
Postal or Zip Code:: 732-0062  
Citizenship Country:: Japan  
Inventor Two Given Name:: Katsumi  
Family Name:: FUJIMOTO  
Postal Address Line One:: 1-3-11-202, Asahi, Minami-ku  
Postal Address Line Two:: Hiroshima-shi  
City:: Hiroshima  
Country:: Japan  
Postal or Zip Code:: 734-0036  
Citizenship Country:: Japan

CORRESPONDENCE INFORMATION

Correspondence Customer Number:: 001444  
Fax One:: 202-737-3528  
Electronic Mail One:: Mail@BrowdyNeimark.com

APPLICATION INFORMATION

Title Line One:: CHONDROGENESIS STIMULATOR  
Total Drawing Sheets:: 6  
Formal Drawings?: Yes  
Docket Number:: KATO=2  
Secrecy Order in Parent Appl.?: No

REPRESENTATIVE INFORMATION

Representative Customer Number:: 1444

CONTINUITY INFORMATION

This application is a:: 371 OF  
> Application One:: PCT/JP00/05590  
Filing Date:: 08-21-2000

PRIOR FOREIGN APPLICATIONS

Foreign Application One:: 232966/1999  
Filing Date:: 08-19-1999  
Country:: Japan

Priority Claimed:: Yes

Source: PrintEFS Version 1.0.1

Source: PrintEFS Version 1.0.1